

Acetyl- and Phospho-Histone H3 (Lys9 Ser10) Blocking Peptide



100 µg
(10 western blots)

Orders ■ 877-616-CELL (2355)
orders@cellsignal.com
Support ■ 877-678-TECH (8324)
info@cellsignal.com
Web ■ www.cellsignal.com

rev. 06/30/17

For Research Use Only. Not For Use In Diagnostic Procedures.

Description: This peptide is used to block Acetyl- and Phospho-Histone H3 (Lys9/Ser10) Antibody #9711 reactivity.

Background: Modulation of chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of DNA wound around eight core histone proteins (two each of H2A, H2B, H3, and H4), is the primary building block of chromatin (1). The amino-terminal tails of core histones undergo various post-translational modifications, including acetylation, phosphorylation, methylation, and ubiquitination (2-5). These modifications occur in response to various stimuli and have a direct effect on the accessibility of chromatin to transcription factors and, therefore, gene expression (6). In most species, histone H2B is primarily acetylated at Lys5, 12, 15, and 20 (4,7). Histone H3 is primarily acetylated at Lys9, 14, 18, 23, 27, and 56. Acetylation of H3 at Lys9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms (2,3). Phosphorylation at Ser10, Ser28, and Thr11 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis (8-10). Phosphorylation at Thr3 of histone H3 is highly conserved among many species and is catalyzed by the kinase haspin. Immunostaining with phospho-specific antibodies in mammalian cells reveals mitotic phosphorylation at Thr3 of H3 in prophase and its dephosphorylation during anaphase (11).

Quality Control: The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry. The peptide blocks Acetyl- and Phospho-Histone H3 (Lys9/Ser10) Antibody #9711 by peptide dot blot.

Applications: Use as a blocking reagent to evaluate the specificity of antibody reactivity in peptide dot blot protocols.

Directions for Use: Use as a blocking reagent to evaluate the specificity of antibody reactivity in peptide dot blot protocols. Recommended antibody dilutions can be found on the relevant product data sheet.

Storage: Supplied in 20 mM potassium phosphate (pH 7.0), 50 mM NaCl, 0.1 mM EDTA, 1 mg/ml BSA, 5% glycerol, and 1% DMSO. Store at -20°C.

Companion Products:

- Acetyl- and Phospho-Histone H3 (Lys9/Ser10) Antibody #9711
- Anti-rabbit IgG, HRP-linked Antibody #7074
- Prestained Protein Marker, Broad Range (Premixed Format) #7720
- Biotinylated Protein Ladder Detection Pack #7727
- 20X LumiGLO® Reagent and 20X Peroxide #7003

Background References:

- (1) Workman, J.L. and Kingston, R.E. (1998) *Annu Rev Biochem* 67, 545-79.
- (2) Hansen, J.C. et al. (1998) *Biochemistry* 37, 17637-41.
- (3) Strahl, B.D. and Allis, C.D. (2000) *Nature* 403, 41-5.
- (4) Cheung, P. et al. (2000) *Cell* 103, 263-71.
- (5) Bernstein, B.E. and Schreiber, S.L. (2002) *Chem Biol* 9, 1167-73.
- (6) Jaskelioff, M. and Peterson, C.L. (2003) *Nat Cell Biol* 5, 395-9.
- (7) Thorne, A.W. et al. (1990) *Eur J Biochem* 193, 701-13.
- (8) Hendzel, M.J. et al. (1997) *Chromosoma* 106, 348-60.
- (9) Goto, H. et al. (1999) *J Biol Chem* 274, 25543-9.
- (10) Preuss, U. et al. (2003) *Nucleic Acids Res* 31, 878-85.
- (11) Dai, J. et al. (2005) *Genes Dev* 19, 472-88.