

Bcl-11B (D6F1) XP® Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP, IF-F, IF-IC, ChIP, ChIP-seq	H M	Endogenous	120-130	Rabbit IgG	#Q9C0K0	64919

Product Usage Information

For optimal ChIP and ChIP-seq results, use 10 µl of antibody and 10 µg of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP® Enzymatic Chromatin IP Kits.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:100
Immunofluorescence (Frozen)	1:100
Immunofluorescence (Immunocytochemistry)	1:200
Chromatin IP	1:50
Chromatin IP-seq	1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #16148.

Specificity/Sensitivity

Bcl-11B (D6F1) XP® Rabbit mAb recognizes endogenous levels of total Bcl-11B protein.

Species predicted to react based on 100% sequence homology

Rat, Monkey

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu391 of human Bcl-11B protein.

Background

Bcl-11B (Ctip2) is a COUP-TF interacting protein that belongs to the C₂H₂ type zinc finger protein family (1). Bcl-11B is highly expressed in the brain and is critical for the development of neurons, as well as other tissues and organs. Bcl-11B also plays an essential role in T cell lineage commitment and maintenance of T cell identity (1-3). Two isoforms of Bcl-11B are found to be encoded by the *BCL11B* gene, possibly through exon-skipping (4). Bcl-11B is a transcription factor which binds to target genes through the 2nd and 3rd zinc-finger domains of exon 4 (3), while also interacting with various protein partners including COUP-TF proteins (1), the NuRD complex (5,6), HDAC1, HDAC2, and SUV39H1 (7). Research studies have shown that mutations and deletion of Bcl-11B contribute to the development of thymic lymphoma in mice and T cell acute lymphoblastic leukemia in humans, indicating a role as a tumor suppressor (4,8). Mechanistic studies have shown that Bcl-11B represses gene expression of the E3 ubiquitin ligase HDM2 in a p53-dependent manner (9).

Background References

1. Avram, D. et al. (2000) *J Biol Chem* 275, 10315-22.
2. Liu, P. et al. (2010) *Immunol Rev* 238, 138-49.
3. Kominami, R. (2012) *Proc Jpn Acad Ser B Phys Biol Sci* 88, 72-87.
4. Wakabayashi, Y. et al. (2003) *Biochem Biophys Res Commun* 301, 598-603.
5. Cismasiu, V.B. et al. (2005) *Oncogene* 24, 6753-64.
6. Topark-Ngarm, A. et al. (2006) *J Biol Chem* 281, 32272-83.
7. Marban, C. et al. (2007) *EMBO J* 26, 412-23.
8. Gutierrez, A. et al. (2011) *Blood* 118, 4169-73.
9. Obata, M. et al. (2012) *Cell Signal* 24, 1047-52.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IF-F:** Immunofluorescence (Frozen) **IF-IC:** Immunofluorescence (Immunocytochemistry) **ChIP:** Chromatin IP **ChIP-seq:** Chromatin IP-seq

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.