

Store at
-20C
#12140**CP110 Antibody**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H Mk	Sensitivity: Endogenous	MW (kDa): 130	Source/Isotype: Rabbit	UniProt ID: #O43303	Entrez-Gene Id: 9738
---------------------------	----------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CP110 Antibody recognizes endogenous levels of total CP110 protein. This antibody cross-reacts with a protein of unknown origin at ~65 kDa.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues around Ser580 of human CP110 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The centrosome is an organelle that plays an important role in the mammalian cell cycle. Prior to the onset of mitosis, the single interphase centrosome duplicates only once, creating a pair of daughter centrosomes that will form the two spindle poles after breakdown of the nuclear envelope. Failure to duplicate or overduplication of the centrosome can result in polyploidy and genomic instability (reviewed in 1). Centrosomal protein of 110 kDa (CP110) is a cyclin-dependent kinase (CDK) substrate that plays a critical role in promoting the duplication of centrosomes and correct spindle formation (2). In addition, CP110 has been shown to interact with calmodulin (CaM) and centrin to regulate the progression through cytokinesis (3), and with Cep97 and Cep290 to regulate the formation of primary cilia (4,5). CP110 expression is induced in G1/S with peak expression during S-phase. Degradation of CP110 is mediated by cyclin F in G2-phase and is required for normal progression into M-phase (6).

Background References

1. Hinchcliffe, E.H. and Sluder, G. (2001) *Genes Dev* 15, 1167-81.
2. Chen, Z. et al. (2002) *Dev Cell* 3, 339-50.
3. Tsang, W.Y. et al. (2006) *Mol Biol Cell* 17, 3423-34.
4. Spektor, A. et al. (2007) *Cell* 130, 678-90.
5. Tsang, W.Y. et al. (2008) *Dev Cell* 15, 187-97.
6. D'Angiolella, V. et al. (2010) *Nature* 466, 138-42.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.