



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at +4C  
#12230

## Histone H3 (D1H2) XP<sup>®</sup> Rabbit mAb (Alexa Fluor<sup>®</sup> 647 Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
IF-IC, FC-FP	H M R Mk	Endogenous	Rabbit IgG	#P68431	8350
<b>Product Usage Information</b>	<b>Application</b>				<b>Dilution</b>
	Immunofluorescence (Immunocytochemistry)				1:200 - 1:800
	Flow Cytometry (Fixed/Permeabilized)				1:50
<b>Storage</b>	Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.				
<b>Specificity/Sensitivity</b>	Histone H3 (D1H2) XP <sup>®</sup> Rabbit mAb detects endogenous levels of total Histone H3 protein, including isoforms H3.1, H3.2, and H3.3. This antibody also detects the Histone H3 variant CENP-A. This antibody does not cross-react with other core histones.				
<b>Species predicted to react based on 100% sequence homology</b>	Hamster, Chicken, D. melanogaster, Xenopus, Zebrafish, Bovine				
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the carboxy terminus of the human histone H3 protein.				
<b>Description</b>	This Cell Signaling Technology antibody is conjugated to Alexa Fluor <sup>®</sup> 647 fluorescent dye and tested in-house for direct flow cytometry and immunofluorescent analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated Histone H3 (D1H2) XP <sup>®</sup> Rabbit mAb #4499.				
<b>Background</b>	Modulation of chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of DNA wound around eight core histone proteins (two each of H2A, H2B, H3, and H4), is the primary building block of chromatin (1). The amino-terminal tails of core histones undergo various posttranslational modifications, including acetylation, phosphorylation, methylation, and ubiquitination (2-5). These modifications occur in response to various stimuli and have a direct effect on the accessibility of chromatin to transcription factors and, therefore, gene expression (6). In most species, histone H2B is primarily acetylated at Lys5, 12, 15, and 20 (4,7). Histone H3 is primarily acetylated at Lys9, 14, 18, 23, 27, and 56. Acetylation of H3 at Lys9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms (2,3). Phosphorylation at Ser10, Ser28, and Thr11 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis (8-10). Phosphorylation at Thr3 of histone H3 is highly conserved among many species and is catalyzed by the kinase haspin. Immunostaining with phospho-specific antibodies in mammalian cells reveals mitotic phosphorylation at Thr3 of H3 in prophase and its dephosphorylation during anaphase (11).				
<b>Background References</b>	<ol style="list-style-type: none"> <li>1. Workman, J.L. and Kingston, R.E. (1998) <i>Annu Rev Biochem</i> 67, 545-79.</li> <li>2. Hansen, J.C. et al. (1998) <i>Biochemistry</i> 37, 17637-41.</li> <li>3. Strahl, B.D. and Allis, C.D. (2000) <i>Nature</i> 403, 41-5.</li> <li>4. Cheung, P. et al. (2000) <i>Cell</i> 103, 263-71.</li> <li>5. Bernstein, B.E. and Schreiber, S.L. (2002) <i>Chem Biol</i> 9, 1167-73.</li> <li>6. Jaskelioff, M. and Peterson, C.L. (2003) <i>Nat Cell Biol</i> 5, 395-9.</li> <li>7. Thorne, A.W. et al. (1990) <i>Eur J Biochem</i> 193, 701-13.</li> <li>8. Hendzel, M.J. et al. (1997) <i>Chromosoma</i> 106, 348-60.</li> <li>9. Goto, H. et al. (1999) <i>J Biol Chem</i> 274, 25543-9.</li> <li>10. Preuss, U. et al. (2003) <i>Nucleic Acids Res</i> 31, 878-85.</li> <li>11. Dai, J. et al. (2005) <i>Genes Dev</i> 19, 472-88.</li> </ol>				

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Applications Key

**IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

## Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Alexa Fluor is a registered trademark of Life Technologies Corporation.

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com).

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.