

Plectin-1 (D6A11) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-IC	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 400-500	Source/Isotype: Rabbit IgG	UniProt ID: #Q15149	Entrez-Gene Id: 5339
--------------------------------------	--------------------------------	-----------------------------------	-----------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting
Immunoprecipitation
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:50
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Plectin-1 (D6A11) Rabbit mAb recognizes endogenous levels of total plectin-1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu2980 of human plectin-1 protein.

Background

Plectin is a large, widely expressed protein that crosslinks the intermediate filament and actin cytoskeleton, mechanically stabilizing cells and tissues. Plectin also plays a role in the regulation of actin dynamics and acts as a scaffold for signaling molecules (1). Plectin is important in the stabilization of hemidesmosomes, crosslinking them to the intermediate filament network. Research studies have shown that mutations in plectin and other genes coding for hemidesmosomal proteins can cause epidermolysis bullosa, a condition manifested by fragile skin and frequent blistering (1,2). Plectin modulates signals to PKC through binding and sequestration of RACK1, the receptor for activated C kinase 1 (3,4). Plectin is also involved in the regulation of cyokeratin architecture and cell stress response (4), signaling through the chemokine receptor CXCR4 (5) and regulation of AMP-activated protein kinase (AMPK) activity and signaling in mouse myotubes (6).

Background References

1. Wiche, G. (1998) *J Cell Sci* 111 (Pt 17), 2477-86.
2. Pfendner, E. et al. (2005) *Exp Dermatol* 14, 241-9.
3. Osmanagic-Myers, S. and Wiche, G. (2004) *J Biol Chem* 279, 18701-10.
4. Osmanagic-Myers, S. et al. (2006) *J Cell Biol* 174, 557-68.
5. Ding, Y. et al. (2008) *Exp Cell Res* 314, 590-602.
6. Gregor, M. et al. (2006) *J Cell Sci* 119, 1864-75.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.