

Store at
-20C
#12292**TEAD1 (D9X2L) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, W-S, IP, IHC-P, IF-IC	Reactivity: H M Mk	Sensitivity: Endogenous	MW (kDa): 50	Source/Isotype: Rabbit IgG	UniProt ID: #P28347	Entrez-Gene Id: 7003
-----------------------------------------------------	------------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting
Simple Western™
Immunoprecipitation
Immunohistochemistry (Paraffin)
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:10 - 1:50
1:100
1:50 - 1:200
1:50 - 1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #95633.

Specificity/Sensitivity

TEAD1 (D9X2L) Rabbit mAb recognizes endogenous levels of total TEAD1 protein.

Species predicted to react based on 100% sequence homology

Rat

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Phe193 of human TEAD1 protein.

Background

The Hippo pathway is an important evolutionarily conserved signaling pathway that controls organ size and tumor suppression by inhibiting cell proliferation and promoting apoptosis (1,2). An integral function of the Hippo pathway is to repress the activity of Yes-associated protein (YAP), a proposed oncogene whose activity is regulated by phosphorylation and subcellular localization (3,4). When the Hippo pathway is turned on, YAP is phosphorylated by LATS1/2 kinase and sequestered in the cytoplasm by 14-3-3 protein binding, rendering YAP inactive. When the Hippo pathway is off, non-phosphorylated YAP translocates to the nucleus where it associates with various transcription factors including members of the transcriptional enhancer factor (TEF) family, also known as the TEA domain (TEAD) family (TEAD1-4) (5,6). Although widely expressed in tissues, the TEAD family proteins have specific tissue and developmental distributions. YAP/TEAD complexes regulate the expression of genes involved in cell proliferation and apoptosis (5).

Background References

1. Pan, D. (2010) *Dev Cell* 19, 491-505.
2. Harvey, K.F. et al. (2003) *Cell* 114, 457-67.
3. Zhao, B. et al. (2010) *Genes Dev* 24, 862-74.
4. Zhao, B. et al. (2008) *Curr Opin Cell Biol* 20, 638-46.
5. Zhao, B. et al. (2008) *Genes Dev* 22, 1962-71.
6. Zhao, B. et al. (2007) *Genes Dev* 21, 2747-61.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **W-S:** Simple Western™ **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin) **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human **M:** Mouse **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SignalStain is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.