

Phospho-Scribble (Ser1220) (D8A2) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M	Endogenous	240	Rabbit IgG	#Q14160-1	23513

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-Scribble (Ser1220) (D8A2) Rabbit mAb recognizes endogenous levels of scribble protein only when phosphorylated at Ser1220.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser1220 of human scribble protein.

Background

Scribble (Scrib) was originally identified in a genetic screen in *Drosophila* along with cell polarity determinants Discs Large (Dlg) and Lethal giant larvae (Lgl). *Drosophila* mutants homozygous for these genes share similar phenotypes, including the loss of apicobasal cell polarity and neoplastic tissue overgrowth. These phenotypic similarities suggest that these three proteins function in a common pathway important for establishing and maintaining apicobasal polarity in epithelial cells (1,2). Scribble contains many leucine-rich repeats and PDZ domains important for localizing scribble to adherens junctions and basolateral regions of mammalian epithelial cells (3). Scribble reportedly binds β-catenin, APC, E-cadherin and the E6 protein from high-risk virus type of HPV through a short motif important for E6-induced cell transformation (4-8). Overexpression of scribble inhibits transformation of rodent epithelial cells by HPV E6/7 proteins (8).

The phosphorylation state of Scribble has been shown to be functionally important, in part by regulating subcellular localization (9). Mass spectrometry studies have identified phosphorylation at Ser1220 as a frequent modification in a variety of cell and tissue types (10-13). The functional significance of this modification remains to be elucidated.

Background References

1. Bilder, D. and Perrimon, N. (2000) *Nature* 403, 676-80.
2. Bilder, D. et al. (2000) *Science* 289, 113-6.
3. Humbert, P.O. et al. (2008) *Oncogene* 27, 6888-907.
4. Sun, Y. et al. (2009) *Mol Biol Cell* 20, 3390-400.
5. Qin, Y. et al. (2005) *J Cell Biol* 171, 1061-71.
6. Navarro, C. et al. (2005) *Oncogene* 24, 4330-9.
7. Takizawa, S. et al. (2006) *Genes Cells* 11, 453-64.
8. Nguyen, M.L. et al. (2003) *J Virol* 77, 6957-64.
9. Yoshihara, K. et al. (2011) *Exp Cell Res* 317, 413-22.
10. Olsen, J.V. et al. (2010) *Sci Signal* 3, ra3.
11. Han, G. et al. (2010) *Electrophoresis* 31, 1080-9.
12. Brill, L.M. et al. (2009) *Cell Stem Cell* 5, 204-13.
13. Wang, Y.T. et al. (2010) *J Proteome Res* 9, 5582-97.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.