

PI3 Kinase Class II α (D3Q5B) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 180	Source/Isotype: Rabbit IgG	UniProt ID: #O00443	Entrez-Gene Id: 5286	
Product Usage Information		Application Western Blotting Immunoprecipitation			Dilution 1:1000 1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Specificity/Sensitivity		PI3 Kinase Class II α (D3Q5B) Rabbit mAb recognizes endogenous levels of total PI3K class II α protein.					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly717 of human PI3K class II α protein.					
Background		Class II phosphatidylinositol 3-kinases (PI3K) contain a C-terminal C2 domain that is unique to the class II isoforms of the PI3K family. This C2 domain mediates protein and phospholipid binding acitivities (1,2). PI3K Class II α generates phosphatidylinositol 3-phosphate (PIP3) and phosphatidylinositol 3,4-bisphosphate (PI(3, 4)P2) from phosphatidylinositol and phosphatidylinositol 4-phosphate (3). PI3K Class II α is located in various intracellular locations such as the trans-Golgi network, endocytic compartments, clathrin-coated vesicles, and nuclear speckles (1,4,5). Research studies have indicated that PI3K Class II α regulates the assembly and distribution of clathrin, resulting in the modulation of clathrin-dependent trafficking and sorting within the trans Golgi network (5,6). PI3K Class II α also mediates translocation of the glucose transporter GLUT4 to the plasma membrane in response to insulin (7). PI3K Class II α has also been shown to regulate neurosecretory granule exocytosis (8) and vascular smooth muscle contraction (9). Unlike other PI3K family members, PI3K Class II α is less sensitive to the PI3K inhibitors wortmannin and LY294002 (3).					
Background Re	eferences	2. Stahelin, R.V. et al. (3. Domin, J. et al. (199 4. Domin, J. et al. (200 5. Gaidarov, I. et al. (2 6. Gaidarov, I. et al. (2 7. Falasca, M. et al. (208 8. Wen, P.J. et al. (2008	2006) J Biol Chem 2 7) Biochem J 326 (F 0) J Biol Chem 275, 001) Mol Cell 7, 443 005) J Biol Chem 28 007) J Biol Chem 28 3) Mol Biol Cell 19, 5	Biol Chem 280, 40766-72. Biol Chem 282, 28226-36.			

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat

dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key H: Human M: Mouse R: Rat Mk: Monkey

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.