



Orders: 877-616-CELL (2355)  
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#12428 Store at +4C

## Phospho-SMAD1 (Ser463/465)/ SMAD5 (Ser463/465)/ SMAD9 (Ser465/467) (D5B10) Rabbit mAb (PE Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

|                               |                             |                                   |                                      |  |  |
|-------------------------------|-----------------------------|-----------------------------------|--------------------------------------|--|--|
| <b>Applications:</b><br>FC-FP | <b>Reactivity:</b><br>H M R | <b>Sensitivity:</b><br>Endogenous | <b>Source/Isotype:</b><br>Rabbit IgG | <b>UniProt ID:</b><br>#Q99717, #Q15797,<br>#O15198 | <b>Entrez-Gene Id:</b><br>4090, 4086, 4093 |
|-------------------------------|-----------------------------|-----------------------------------|--------------------------------------|--|--|

|   |  |                         |
|---|--|-------------------------|
| <b>Product Usage Information</b>                                  | <b>Application</b><br>Flow Cytometry (Fixed/Permeabilized)   | <b>Dilution</b><br>1:50 |
| <b>Storage</b>  | Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.   |                         |
| <b>Specificity/Sensitivity</b>                                    | Phospho-SMAD1 (Ser463/465)/ SMAD5 (Ser463/465)/ SMAD9 (Ser465/467) (D5B10) Rabbit mAb (PE Conjugate) recognizes endogenous levels of SMAD1 and SMAD5 protein when phosphorylated at Ser463/465 and SMAD9 (SMAD8) protein when phosphorylated at Ser465/467.  |                         |
| <b>Species predicted to react based on 100% sequence homology</b> | Monkey   |                         |
| <b>Source / Purification</b>                                      | Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser463/465 of human SMAD1 and SMAD5 protein.   |                         |
| <b>Description</b>  | This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-SMAD1 (Ser463/465)/ SMAD5 (Ser463/465)/ SMAD9 (Ser465/467) (D5B10) Rabbit mAb #13820.   |                         |
| <b>Background</b>   | Bone morphogenetic proteins (BMPs) constitute a large family of signaling molecules that regulate a wide range of critical processes including morphogenesis, cell-fate determination, proliferation, differentiation, and apoptosis (1,2). BMP receptors are members of the TGF- $\beta$ superfamily of Ser/Thr kinase receptors. Ligand binding induces multimerization, autophosphorylation, and activation of these receptors (3-5). They subsequently phosphorylate SMAD1 at Ser463 and Ser465 in the carboxy-terminal motif SSXS, as well as SMAD5 and SMAD9 (SMAD8) at their corresponding sites. These phosphorylated SMADs dimerize with the coactivating SMAD4 and translocate to the nucleus, where they regulate the transcription of target genes (5). MAP kinases and CDKs 8 and 9 are also reported to phosphorylate residues in the linker region of SMAD1, including Ser206. Phosphorylation of SMAD1 at Ser206 recruits Smurf1 to the linker region and leads to the degradation of SMAD1 (6). Phosphorylation at this site also promotes SMAD1 transcriptional activity by recruiting YAP to the linker region (7). |                         |
| <b>Background References</b>                                      | <ol style="list-style-type: none"> <li>Hogan, B.L. (1996) <i>Genes Dev</i> 10, 1580-94.</li> <li>Hoodless, P.A. et al. (1996) <i>Cell</i> 85, 489-500.</li> <li>Klemm, J.D. et al. (1998) <i>Annu Rev Immunol</i> 16, 569-92.</li> <li>Kretzschmar, M. et al. (1997) <i>Genes Dev</i> 11, 984-95.</li> <li>Whitman, M. (1998) <i>Genes Dev</i> 12, 2445-62.</li> <li>Sapkota, G. et al. (2007) <i>Mol Cell</i> 25, 441-54.</li> <li>Alarcón, C. et al. (2009) <i>Cell</i> 139, 757-69.</li> </ol>  |                         |

**Species Reactivity** Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Applications Key** **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

**Cross-Reactivity Key** **H:** Human **M:** Mouse **R:** Rat

**Trademarks and Patents** Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.