

Store at
-20C
#12511**FGF Receptor 1 (D8E4) XP[®] Rabbit mAb (Biotinylated)**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M Mk	Endogenous	92, 120, 145	Rabbit IgG	#P11362	2260

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
Storage	Supplied in 140 mM NaCl, 3 mM KCl, 10 mM sodium phosphate (pH 7.4) dibasic, 2 mM potassium phosphate monobasic, 2 mg/mL BSA, and 50% glycerol. Store at -20°C. <i>Do not aliquot the antibody.</i>	
Specificity/Sensitivity	FGF Receptor 1 (D8E4) XP [®] Rabbit mAb (Biotinylated) recognizes endogenous levels of total FGF receptor 1 protein. This antibody does not cross-react with other FGF receptor family members.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a recombinant protein specific to the carboxy terminus of human FGF receptor 1 protein.	
Description	This Cell Signaling Technology antibody is conjugated to biotin under optimal conditions. The biotinylated antibody is expected to exhibit the same species cross-reactivity as the unconjugated FGF Receptor 1 (D8E4) XP [®] Rabbit mAb #9740.	
Background	Fibroblast growth factors (FGFs) produce mitogenic and angiogenic effects in target cells by signaling through cell surface receptor tyrosine kinases. There are four members of the FGF receptor family: FGFR1 (flg), FGFR2 (bek, KGFR), FGFR3, and FGFR4. Each receptor contains an extracellular ligand-binding domain, a transmembrane domain, and a cytoplasmic kinase domain (1). Following ligand binding and dimerization, the receptors are phosphorylated at specific tyrosine residues (2). Seven tyrosine residues in the cytoplasmic tail of FGFR1 can be phosphorylated: Tyr463, 583, 585, 653, 654, 730, and 766. Tyr653 and Tyr654 are important for catalytic activity of activated FGFR and are essential for signaling (3). The other phosphorylated tyrosine residues may provide docking sites for downstream signaling components, such as Crk and PLCγ (4,5).	
Background References	<ol style="list-style-type: none"> 1. Powers, C.J. et al. (2000) <i>Endocr Relat Cancer</i> 7, 165-97. 2. Reilly, J.F. et al. (2000) <i>J Biol Chem</i> 275, 7771-8. 3. Mohammadi, M. et al. (1996) <i>Mol Cell Biol</i> 16, 977-89. 4. Mohammadi, M. et al. (1991) <i>Mol Cell Biol</i> 11, 5068-78. 5. Larsson, H. et al. (1999) <i>J Biol Chem</i> 274, 25726-34. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key	W: Western Blotting	
Cross-Reactivity Key	H: Human M: Mouse Mk: Monkey	
Trademarks and Patents	<p>Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.</p> <p>XP is a registered trademark of Cell Signaling Technology, Inc.</p> <p>All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.</p>	
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.	

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.