

Store at
-20C
#12619**CTR9 (D1Z4F) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R	Endogenous	140	Rabbit IgG	#Q6PD62	9646

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CTR9 (D1Z4F) Rabbit mAb recognizes endogenous levels of total CTR9 protein.

Species predicted to react based on 100% sequence homology

Hamster, Chicken, Xenopus, Zebrafish, Bovine, Dog, Pig

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human CTR9 protein.

Background

The PAF (RNA polymerase II (RNAPII) associated factor) complex was initially identified in yeast and is comprised of subunits PAF1, Leo1, Ctr9, Cdc73, RTF1 and Ski8 (1,2). The PAF complex plays an important role in transcription initiation and elongation by RNAPII by regulating the establishment of proper histone modifications such as histone H2B ubiquitination and the recruitment of the histone chaperone FACT (facilitates chromatin transcription) (3-5). The PAF complex also plays a role in mRNA processing and maturation by interacting with and recruiting the cleavage and polyadenylation specificity factor and cleavage stimulation factor complexes via the Cdc73 subunit (6,7). In addition, the Ski8 subunit of the PAF complex is part of the hSKI complex that regulates RNA surveillance, suggesting an important function of the complex in coordinating events associated with proper RNA maturation during transcription (1,5).

Background References

1. Jaehning, J.A. (2010) *Biochim Biophys Acta* 1799, 379-88.
2. Chaudhary, K. et al. (2007) *Oncogene* 26, 7499-507.
3. Pavri, R. et al. (2006) *Cell* 125, 703-17.
4. Rozenblatt-Rosen, O. et al. (2005) *Mol Cell Biol* 25, 612-20.
5. Zhu, B. et al. (2005) *Genes Dev* 19, 1668-73.
6. Rozenblatt-Rosen, O. et al. (2009) *Proc Natl Acad Sci U S A* 106, 755-60.
7. Farber, L.J. et al. (2010) *Mol Carcinog* 49, 215-23.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.