

**Atlastin-1 (D2E6) Rabbit mAb**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W	<b>Reactivity:</b> H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 55	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #Q8WXF7	<b>Entrez-Gene Id:</b> 51062
---------------------------	-----------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	---------------------------------

**Product Usage Information****Application**

Western Blotting

**Dilution**

1:1000

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

Atlastin-1 (D2E6) Rabbit mAb recognizes endogenous levels of total atlastin-1 protein. This antibody does not cross-react with atlastin-2 or atlastin-3. This antibody may cross-react with a protein of unknown origin at 100 kDa.

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human atlastin-1 protein.

**Background**

Atlastin proteins are highly conserved members of the dynamin superfamily of membrane GTPases that are involved in the formation of vesicles for the endocytotic and secretory processes (1). Atlastins are required to establish and maintain the morphology of the tubular endoplasmic reticulum (ER) and are therefore important in ER function (2). GTP hydrolysis and dimerization are required for atlastin-dependent ER membrane fusion (3).

Atlastin GTPase 1 (ATL1) is primarily expressed in brain, while the related atlastin 2 and atlastin 3 proteins are ubiquitously expressed (4). Mutations in the atlastin 1 gene SPG3A and the ER defects that result are thought to cause one form of hereditary spastic paraplegia (HSP), a group of heterogeneous neurological disorders characterized by severe progressive spasticity of the lower limbs (5,6).

**Background References**

1. McNiven, M.A. et al. (2000) *Trends Biochem Sci* 25, 115-20.
2. Park, S.H. and Blackstone, C. (2010) *EMBO Rep* 11, 515-21.
3. Moss, T.J. et al. (2011) *Proc Natl Acad Sci U S A* 108, 11133-8.
4. Rismanchi, N. et al. (2008) *Hum Mol Genet* 17, 1591-604.
5. Salinas, S. et al. (2008) *Lancet Neurol* 7, 1127-38.
6. Hu, J. et al. (2009) *Cell* 138, 549-61.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting

**Cross-Reactivity Key**

**H:** Human **M:** Mouse **R:** Rat

**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.