

NXF1 (D5X4G) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures

Applications: W	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 70	Source/Isotype: Rabbit IgG	UniProt ID: #Q9UBU9	Entrez-Gene Id: 10482
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		NXF1 (D5X4G) Rabbit mAb recognizes endogenous levels of total NXF1 protein.				
Species predicte based on 100% homology		Hamster, Bovine, Dog,	Pig, Guinea Pig			
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gln211 of human NXF1 protein.				
Background		mRNA export is a process that is tightly coupled to mRNA splicing (1-4). Splicing and packaging of mRNAs in the form of an mRNA-protein complex (mRNP) leads to the recruitment of the mRNA export adaptor THOC4/ALY, via its interaction with the splicing factor UAP56, forming a large complex termed the transcription-export complex (TREX) (1,2,5). THOC4/ALY then directly interacts with NXF1/TAP, a part of the heterodimer that targets the mRNP to the nuclear pore complex, resulting in the shuttling of mRNP out of the nucleus and into the cytoplasm (1-3,6).				
Background References		 Katahira, J. (2012) <i>Biochim Biophys Acta</i> 1819, 507-13. Siddiqui, N. and Borden, K.L. (2012) Wiley Interdiscip Rev RNA 3, 13-25. Reed, R. and Magni, K. (2001) <i>Nat Cell Biol</i> 3, E201-4. Masuda, S. et al. (2005) <i>Genes Dev</i> 19, 1512-7. Luo, M.L. et al. (2001) <i>Nature</i> 413, 644-7. Kang, Y. and Cullen, B.R. (1999) <i>Genes Dev</i> 13, 1126-39. 				
Species Reactiv	itv	Species reactivity is de	etermined by testing	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Ke	у	W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat Mk: Monkey				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				

more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.