## Phospho-Gab1 (Tyr659) Antibody





Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	<b>MW (kDa):</b> 110	Source/Isotype: Rabbit	<b>UniProt ID:</b> #Q13480	Entrez-Gene Id: 2549	
Product Usage Information		<b>Application</b> Western Blotting Immunoprecipitation			<b>Dilution</b> 1:1000 1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity/Sens	itivity	Phosho-Gab1 (Tyr659) Antibody recognizes endogenous levels of Gab1 protein only when phosphorylated at Tyr659.					
Species predicto based on 100% homology	ed to react sequence	Mouse, Rat					
Source / Purifica	ation	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr659 of human Gab1 protein. Antibodies are purified by protein A and peptide affinity chromatography.					
Background		The Grb-associated bir receptor tyrosine kinas antigen receptors. Upo membrane where it is phosphorylation sites binding to p85 PI3 kina association to PLCy (8). activation of the protei	nder (Gab) family is ses (RTKs) such as E on stimulation of R <sup>-</sup> phosphorylated an of Gab1 protein ha ase (6,7). Phosphor . Phosphorylation o n tyrosine phospha	a family of adaptor prot GFR, HGFR, insulin rece FKs by their cognate liga d functions as a scaffolo ve been identified (5). Pł ylation of Gab1 at Tyr30 of Tyr627 and Tyr659 is r atase SHP2 (6,9).	eins recruited by a ptor, cytokine recep nd, Gab is recruited (1-4). Multiple tyro nosphorylation of T 7, Tyr373 and Tyr40 equired for Gab1 b	wide variety of otor and B cell d to the plasma osine yr472 regulates its 07 modulates its inding to and	
Background References 1. Holgado-Madruga, M. et al. (1996) Nature 379, 560-564.   2. Weidner, K.M. et al. (1996) Nature 384, 173-176.   3. Takahashi-Tezuka, M. et al. (1998) Mol. Cell. Biol. 18, 4109   4. Ingham, R.J. et al. (2001) J Biol Chem 276, 12257-65.   5. Lehr, S. et al. (1999) Biochemistry 38, 151-159.   6. Rocchi, S. et al. (1998) Mol. Endocrinol. 12, 914-923.   7. Yu, C.F. et al. (2001) J Biol Chem 276, 32552-8.   8. Gual, P. et al. (2000) Oncogene 19, 1509-18.   9. Cunnick, J.M. et al. (2001) J Biol Chem 276, 24380-7.				<i>ure</i> 379, 560-564. 173-176. <i>Cell. Biol.</i> 18, 4109-4117. 6, 12257-65. 51-159. 12, 914-923. 552-8. 9-18. 76, 24380-7.			
Species Reactiv	ity	Species reactivity is de	termined by testing	g in at least one approve	d application (e.g.,	western blot).	
Western Blot Bu	uffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Ke	У	W: Western Blotting IP	: Immunoprecipita	tion			
Cross-Reactivity	у Кеу	H: Human					
Trademarks and	d Patents	Cell Signaling Technolo	ogy is a trademark	of Cell Signaling Techno	logy, Inc.		
		All other trademarks a more information.	re the property of t	heir respective owners.	Visit cellsignal.com	/trademarks for	
<b>Limited Uses</b> Except as otherwise expressly agreed in a writing signed by a legally authorized return the following terms apply to Products provided by CST, its affiliates or its distribut					ally authorized repr es or its distributors	resentative of CST, 5. Any Customer's	

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.