



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at +4C  
#12777

## FGF Receptor 1 (D8E4) XP<sup>®</sup> Rabbit mAb (PE Conjugate)

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
FC-FP	H M R Mk	Endogenous	Rabbit IgG	#P11362	2260

### Product Usage Information

#### Application

Flow Cytometry (Fixed/Permeabilized)

#### Dilution

1:50

### Storage

Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.

### Specificity/Sensitivity

FGF Receptor 1 (D8E4) XP<sup>®</sup> Rabbit mAb (PE Conjugate) detects endogenous levels of total FGF receptor 1 protein. This antibody does not cross-react with other FGF receptor family members.

### Source / Purification

Monoclonal antibody is produced by immunizing animals with a recombinant protein specific to the carboxy terminus of human FGF receptor 1 protein.

### Description

This Cell Signaling Technology antibody is conjugated to phycoerythrin (PE) and tested in-house for direct flow cytometry analysis in human cells. The antibody is expected to exhibit the same species cross-reactivity as the unconjugated FGF Receptor 1 (D8E4) XP<sup>®</sup> Rabbit mAb #9740.

### Background

Fibroblast growth factors (FGFs) produce mitogenic and angiogenic effects in target cells by signaling through cell surface receptor tyrosine kinases. There are four members of the FGF receptor family: FGFR1 (flg), FGFR2 (bek, KGFR), FGFR3, and FGFR4. Each receptor contains an extracellular ligand-binding domain, a transmembrane domain, and a cytoplasmic kinase domain (1). Following ligand binding and dimerization, the receptors are phosphorylated at specific tyrosine residues (2). Seven tyrosine residues in the cytoplasmic tail of FGFR1 can be phosphorylated: Tyr463, 583, 585, 653, 654, 730, and 766. Tyr653 and Tyr654 are important for catalytic activity of activated FGFR and are essential for signaling (3). The other phosphorylated tyrosine residues may provide docking sites for downstream signaling components, such as Crk and PLCγ (4,5).

### Background References

1. Powers, C.J. et al. (2000) *Endocr Relat Cancer* 7, 165-97.
2. Reilly, J.F. et al. (2000) *J Biol Chem* 275, 7771-8.
3. Mohammadi, M. et al. (1996) *Mol Cell Biol* 16, 977-89.
4. Mohammadi, M. et al. (1991) *Mol Cell Biol* 11, 5068-78.
5. Larsson, H. et al. (1999) *J Biol Chem* 274, 25726-34.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Applications Key

**FC-FP:** Flow Cytometry (Fixed/Permeabilized)

### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.