

## Gelsolin (D9W8Y) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IHC-P	Reactivity: H M R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 83	<b>Source/Isotype:</b> Rabbit IgG	UniProt ID: #P06396	Entrez-Gene Id: 2934
Product Usage Information		Application         Western Blotting         Immunohistochemistry (Paraffin)			<b>Dilution</b> 1:1000 1:600	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.  For a carrier free (BSA and azide free) version of this product see product #57376.				
Specificity/Sensitivity		Gelsolin (D9W8Y) Rabbit mAb recognizes endogenous levels of total gelsolin protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val377 of human gelsolin protein.				
Background		Gelsolin (actin-depolymerizing factor, ADF, AGEL, Brevin) is an 83 kDa protein that shares structural and functional homology to villin and adseverin/scinderin (1,2). Gelsolin plays an important role in actin filament assembly by capping and severing actin proteins in a Ca <sup>2+</sup> -dependent manner (3,4). Gelsolin is important for cellular events (e.g., membrane ruffling, chemotaxis, ciliogenesis) that require cytoskeletal remodeling (3). Accordingly, cells from gelsolin knockout mice exhibit motility defects, including a failure to ruffle in response to growth factor stimulation (5,6). In humans, defects in gelsolin have been linked to amyloidosis type 5 (AMYL5), a hereditary disease characterized by cranial neuropathy, which appears to result from gelsolin amyloid deposition (7).				
Background Re	ferences	1. Vitale, M.L. et al. (1991) <i>J Cell Biol</i> 113, 1057-67. 2. Lueck, A. et al. (1998) <i>J Cell Sci</i> 111 ( Pt 24), 3633-43. 3. Sun, H.Q. et al. (1999) <i>J Biol Chem</i> 274, 33179-82. 4. Li, G.H. et al. (2010) <i>Med Res Rev</i> , 32, 999-1025. 5. Azuma, T. et al. (1998) <i>EMBO J</i> 17, 1362-70. 6. Lu, M. et al. (1997) <i>J Cell Biol</i> 138, 1279-87. 7. Haltia, M. et al. (1990) <i>Biochem Biophys Res Commun</i> 167, 927-32.				

**Species Reactivity** Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting IHC-P: Immunohistochemistry (Paraffin)

Cross-Reactivity Key H: Human M: Mouse R: Rat Mk: Monkey

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for a constant of their respective owners are the property of their respective owners.

more information.

**Limited Uses** Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST,

the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.