3024

#

Phospho-CDCP1 (Tyr806) Antibody



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 140	Source/Isotype: Rabbit	UniProt ID: #Q9H5V8	Entrez-Gene Id: 64866		
Product Usage Information		Application Western Blotting			Dilution 1:1000			
Storage		- Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.						
Specificity/Sen	sitivity	Phospho-CDCP1 (Tyr806) Antibody recognizes endogenous levels of CDCP1 protein only when phosphorylated at Tyr806. This antibody does not cross-react with CDCP1 phosphorylated at Tyr707, Tyr734, or Tyr743, but may cross-react with other phospho-tyrosine containing proteins.						
Species predict based on 100% homology		Mouse, Rat						
Source / Purific	cation	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr806 of human CDCP1 protein. Antibodies are purified by protein A and peptide affinity chromatography.						
Background		CUB domain-containing protein 1 (CDCP1, SIMA135) is a putative stem cell marker shown in research studies to be highly expressed in some human cancer cells and in both typical and atypical (cancerous) colons (1). Expression of CDCP1 may be epigenetically regulated, as methylation of promoter CpG sequences results in decreased CDCP1 expression (2). The corresponding <i>CDCP1</i> gene encodes a glycoprotein that acts as a complex, multidomain transmembrane antigen. CDCP1 has three extracellular CUB domains that may be involved in cell adhesion or extracellular matrix interactions (1,3). Src-family kinases may phosphorylate CDCP1 at five tyrosine residues within its cytoplasmic domain to provide a potential binding site for SH2 domain-containing proteins (3). CDCP1 is a putative hematopoietic stem cell marker (4,5).						
Background Re	eferences	1. Scherl-Mostageer, M. et al. (2001) <i>Oncogene</i> 20, 4402-8. 2. Ikeda, J.I. et al. (2006) <i>J. Pathol.</i> 210, 75-84. 3. Hooper, J.D. et al. (2003) <i>Oncogene</i> 22, 1783-94. 4. Conze, T. et al. (2003) <i>Ann. N.Y. Acad. Sci.</i> 996, 222-226. 5. Bühring, H.J. et al. (2004) <i>Stem Cells</i> 22, 334-343.						
Species Reactiv	vity	Species reactivity is de	termined by testin	g in at least one approve	ed application (e.g.,	western blot).		
Western Blot B	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting						
Cross-Reactivit	ty Key	H: Human						
Trademarks ar	nd Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		All other trademarks a more information.	other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for re information.					
Limited Uses		the following terms an terms and conditions	se expressly agreed in a writing signed by a legally authorized representative of CST, ns apply to Products provided by CST, its affiliates or its distributors. Any Customer's ons that are in addition to, or different from, those contained herein, unless ed in writing by a legally authorized representative of CST, are rejected and are of no					

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.