

PSMB7 (E1L5H) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M Mk	Endogenous	28, 30	Rabbit IgG	#Q99436	5695

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

PSMB7 (E1L5H) Rabbit mAb recognizes endogenous levels of total PSMB7 protein. This antibody reacts with precursor and mature forms of PSMB7, but does not cross-react with PSMB10.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys237 of human PSMB7 protein.

Background

The 26S proteasome is a highly abundant proteolytic complex involved in the degradation of ubiquitinated substrate proteins. It consists largely of two sub-complexes, the 20S catalytic core particle (CP) and the 19S/PA700 regulatory particle (RP) that can cap either end of the CP. The CP consists of two stacked heteroheptameric β -rings (β_{1-7}) that contain three catalytic β -subunits and are flanked on either side by two heteroheptameric α -rings (α_{1-7}). The RP includes a base and a lid, each having multiple subunits. The base, in part, is composed of a heterohexameric ring of ATPase subunits belonging to the AAA (ATPases Associated with diverse cellular Activities) family. The ATPase subunits function to unfold the substrate and open the gate formed by the α -subunits, thus exposing the unfolded substrate to the catalytic β -subunits. The lid consists of ubiquitin receptors and DUBs that function in recruitment of ubiquitinated substrates and modification of ubiquitin chain topology (1,2). Other modulators of proteasome activity, such as PA28/11S REG, can also bind to the end of the 20S CP and activate it (1,2).

The core particle exhibits three distinct enzymatic activities, each catalyzed by a separate protein subunit. The constitutively expressed PSMB5, PSMB7, and PSMB6 subunits provide chymotrypsin-like, trypsin-like, and caspase-like activities, respectively. These catalytic subunits belong to the amino-terminal nucleophile (Ntn) hydrolase family and are characterized by a single-residue active site. The catalytic β -subunits are synthesized with amino-terminal propeptides, which are removed at the final step of proteasome biogenesis to expose the catalytic threonine residues (3). In immune cells involved in antigen presentation, the constitutively expressed PSMB6, PSMB7, and PSMB5 subunits are replaced by three highly homologous, induced β -subunits to form the immunoproteasome (4,5). PSMB7 is downregulated at the protein level by IFN- γ and replaced by PSMB10 to remodel the proteolytic specificity of the proteasome for more appropriate immunological processing of endogenous antigens (6). Research studies show that PSMB7 expression is upregulated in human colon adenocarcinomas and suggest that high PSMB7 expression may serve as a potential prognostic marker in breast cancer (7,8).

Background References

1. Finley, D. (2009) *Annu Rev Biochem* 78, 477-513.
2. Lee, M.J. et al. (2011) *Mol Cell Proteomics* 10, R110.003871.
3. Stringer, J.R. et al. (1977) *J Virol* 21, 889-901.
4. Boes, B. et al. (1994) *J Exp Med* 179, 901-9.
5. Cardozo, C. and Kohanski, R.A. (1998) *J Biol Chem* 273, 16764-70.
6. Hisamatsu, H. et al. (1996) *J Exp Med* 183, 1807-16.
7. Rho, J.H. et al. (2008) *J Proteome Res* 7, 2959-72.
8. Munkácsy, G. et al. (2010) *Br J Cancer* 102, 361-8.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.