

LIMD1 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M	Sensitivity: Endogenous	MW (kDa): 72	Source/Isotype: Rabbit	UniProt ID: #Q9UGP4	Entrez-Gene Id: 8994
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM so 20°C. Do not aliquot t		s), 150 mM NaCl, 100 μg	/ml BSA and 50% g	lycerol. Store at –
Specificity/Sensitivity		LIMD1 Antibody recognizes endogenous levels of total LIMD1 protein. This antibody cross-reacts with a protein of unknown origin at \sim 45 kDa.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human LIMD1 protein. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		that belongs to the Aj mediate protein-prote Research studies indic E2F mediated transcri phosphorylation state Von Hippel-Lindau (VH initiation of HIF-1α ub as a tumor suppresso	uba family of LIM dein interactions and cate that LIMD1 is in ption (2) and negate (3,4). Additional stall) protein and the iquitination and derincludes the down	is a putative tumor supomain containing prote typically contain a pair involved in numerous celive regulation of the Hipudies identify LIMD1 as hydroxylase PHD1 to a production (5). Research regulation of the proteers (7), and altered subcers (7), and altered subcers (7), and altered subcers (7).	ins. LIM domain co of distinct zinc fing Ilular processes, ind opo pathway throug a hypoxia regulato orotein complex tha evidence supportin in in 80% of lung ca	ntaining proteins er domains (1). Eluding inhibition of gh influence on YAP or as it recruits the est promotes g the role of LIMD1 encers (6), loss of
Background References		1. Schmeichel, K.L. and Beckerle, M.C. (1994) <i>Cell</i> 79, 211-9. 2. Sharp, T.V. et al. (2004) <i>Proc Natl Acad Sci U S A</i> 101, 16531-6. 3. Sun, G. and Irvine, K.D. (2013) <i>Sci Signal</i> 6, ra81. 4. Reddy, B.V. and Irvine, K.D. (2013) <i>Dev Cell</i> 24, 459-71. 5. Foxler, D.E. et al. (2012) <i>Nat Cell Biol</i> 14, 201-8. 6. Sharp, T.V. et al. (2008) <i>Proc Natl Acad Sci U S A</i> 105, 19932-7. 7. Ghosh, S. et al. (2010) <i>Mol Cancer</i> 9, 58. 8. Spendlove, I. et al. (2008) <i>Int J Cancer</i> 123, 2247-53.				
Species Reactivit	tv	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.	western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human M: Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.