

Basigin/EMMPRIN (E1S1V) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IHC-P	H	Endogenous	38-58	Rabbit IgG	#P35613	682

Product Usage Information**Application**

Western Blotting
Immunohistochemistry (Paraffin)

Dilution

1:1000
1:200 - 1:800

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #95203.

Specificity/Sensitivity

Basigin/EMMPRIN (E1S1V) Rabbit mAb recognizes endogenous levels of total EMMPRIN protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala254 of human EMMPRIN protein.

Background

Basigin (EMMPRIN, CD147) is a type I integral membrane receptor protein belonging to the immunoglobulin superfamily (1). Basigin is a glycosylated protein with four known isoforms, of which isoform 2 is the most abundantly expressed (2). Multiple functions have been ascribed to Basigin; foremost among these is stimulating the secretion of extracellular matrix metalloproteinases by adjacent fibroblasts, a function which has been implicated in promoting tumor progression (2-4). Research studies have shown that Basigin is overexpressed by many tumor cells, and its expression level may correlate with tumor malignancy (5,6). A recent study identified the *BASIGIN* gene as a regulatory target of Slug, suggesting a role for Basigin in the process of epithelial-mesenchymal transition (7). Basigin has also been identified as a marker for a subset of highly suppressive regulatory T cells (8), and as an obligate receptor for the malarial parasite *Plasmodium falciparum* on human erythrocytes (9).

Background References

1. Biswas, C. et al. (1995) *Cancer Res* 55, 434-9.
2. Liao, C.G. et al. (2011) *Mol Cell Biol* 31, 2591-604.
3. Sweeny, L. et al. (2012) *Exp Cell Res* 318, 1788-98.
4. Lescaille, G. et al. (2012) *BMC Cancer* 12, 115.
5. Piao, S. et al. (2012) *J Cancer Res Clin Oncol* 138, 627-35.
6. Bi, X.C. et al. (2011) *Clin Invest Med* 34, E358.
7. Wu, J. et al. (2011) *Oncogene* 30, 4410-27.
8. Solstad, T. et al. (2011) *Blood* 118, 5141-51.
9. Crosnier, C. et al. (2011) *Nature* 480, 534-7.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IHC-P:** Immunohistochemistry (Paraffin)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SignalStain is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.