3404

Phospho-Rad17 (Ser635) (D6K9P) Rabbit



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 80	Source/Isotype: Rabbit IgG	UniProt ID: #075943	Entrez-Gene Id: 5884		
Product Usage Information	9	Application Western Blotting			Dilution 1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Ser	pecificity/Sensitivity Phospho-Rad17 (Ser635) (D6K9P) Rabbit mAb recognizes endogenous levels of Rad17 protein when phosphorylated at Ser635.				7 protein only			
Source / Purifi	cation	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser635 of human Rad17 protein.						
Background		The human checkpoint protein Rad17 and its fission and budding yeast orthologues (<i>Schizosaccharomyces pombe</i> Rad17 and <i>Saccharomyces cerevisiae</i> Rad24, respectively) are involved in the activation of checkpoint signals in response to DNA damage or disruption of DNA synthesis (1-4). Treatment of human cells with genotoxic agents induces ATM/ATR-dependent phosphorylation of Rad17 at Ser635 and Ser645. Rad17 phosphorylation is a critical early event during checkpoint signaling in DNA-damaged cells (5-7).						
Background R	eferences	1. Griffiths, D.J. et al. (2. Li, L. et al. (1999) <i>O</i> 3. Bao, S. et al. (1998) 4. von Deimling, F. et 5. Bao, S. et al. (2001) 6. Post, S. et al. (2001) 7. Wang, X. et al. (200	ncogene 18, 1689-9 Cell Growth Differ 9 al. (1999) Hum Gen Nature 411, 969-74 Proc Natl Acad Sci	9.), 961-7. <i>et</i> 105, 17-27. <i>U S A</i> 98, 13102-7.				
Species Reacti	vity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot E	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting						
Cross-Reactivi	ty Key	H: Human						
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses		the following terms a terms and conditions	rise expressly agreed in a writing signed by a legally authorized representative of CST, ms apply to Products provided by CST, its affiliates or its distributors. Any Customer's cions that are in addition to, or different from, those contained herein, unless ted in writing by a legally authorized representative of CST, are rejected and are of no					
		approved, cleared, or purpose. Customer sl any manner that conf Customer as the end-	licensed by the FDA nall not use any Pro licts with its labeling user and solely for	se Only or a similar labe or other regulatory for duct for any diagnostic o statement. Products so research and developme urposes, or any purchase	eign or domestic er or therapeutic purp old or licensed by C ent uses. Any use of	itity, for any ose, or otherwise in ST are provided for Product for		

component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.