

## VAMP2 (D6O1A) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only, Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IP, IHC-P, IF-F, IF- IC	Reactivity: H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 13	Source/Isotype: Rabbit	<b>UniProt ID:</b> #P63027	Entrez-Gene Id: 6844
Product Usage Information		Application			Dilution	
		Western Blotting			1:1000	)
		Immunoprecipitation			1:50	
		Immunohistochemist	-			0 - 1:6400
		Immunofluorescence	` ,		1:2000	)
		Immunofluorescence	(Immunocytochem	istry)	1:250	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
		For a carrier free (BSA and azide free) version of this product see product #50423.				
Specificity/Sensitivity		VAMP2 (D6O1A) Rabbit mAb recognizes endogenous levels of total VAMP2 protein. This antibody does not cross-react with VAMP1 protein and recognizes VAMP2 protein by IF-IC analysis in mouse and rat cells.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the amino terminus of human VAMP2 protein.				
Background		Vesicle-associated membrane protein 2 (VAMP2, also called synaptobrevin) is part of the R-soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complex (1). The SNARE complex is involved in vesicular transport and membrane fusion, a process regulated by calcium (2). In neurons, VAMP2 is predominantly inserted in presynaptic vesicle membranes. Assembly of VAMP2 with the plasma membrane SNAREs syntaxin 1 and SNAP25 is a key event necessary for membrane fusion and neurotransmitter release (2). In addition to this important function, VAMP2 is also involved in granule exocytosis in neutrophils (3) and release of bioactive peptides from cardiac myocytes (4) and juxtaglomerular cells (5).				
Background References		<ol> <li>Baumert, M. et al. (1989) EMBO J 8, 379-84.</li> <li>Sudhof, T.C. (2004) Annu Rev Neurosci 27, 509-47.</li> <li>Mollinedo, F. et al. (2003) J Immunol 170, 1034-42.</li> <li>Ferlito, M. et al. (2010) J Mol Cell Cardiol 49, 791-800.</li> <li>Mendez, M. et al. (2011) J Biol Chem 286, 28608-18.</li> </ol>				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				

**Applications Key** 

**W:** Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin) **IF-F:** Immunofluorescence (Frozen) **IF-IC:** Immunofluorescence (Immunocytochemistry)

**Cross-Reactivity Key** 

H: Human M: Mouse R: Rat

**Trademarks and Patents** 

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Alexa Fluor is a registered trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

## **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.