

MW (kDa): **Applications:** Reactivity: Sensitivity: Source/Isotype: UniProt ID: Entrez-Gene Id: W, IP, ChIP HMRMk Rabbit IgG #Q08999 Endogenous 130 5934 For optimal ChIP results, use 10 μ l of antibody and 10 μ g of chromatin (approximately 4 x 10⁶ cells) per Product Usage IP. This antibody has been validated using SimpleChIP[®] Enzymatic Chromatin IP Kits. Information Application Dilution 1:1000 Western Blotting 1:100 Immunoprecipitation Chromatin IP 1:50 Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than Storage 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody. RBL2 (D9T7M) Rabbit mAb recognizes endogenous levels of total RBL2 protein. This antibody does not Specificity/Sensitivity cross-react with the Rb or RBL1 proteins. Source / Purification Monoclonal antibody is produced by immunizing animals with recombinant protein corresponding to human RBL2 protein. Background The retinoblastoma (Rb) tumor suppressor family includes the retinoblastoma protein Rb (p105), retinoblastoma-like protein 1 (RBL1, p107), and retinoblastoma-like protein 2 (RBL2, p130). These Rb family proteins are referred to as 'pocket proteins' because they contain a conserved binding pocket region that interacts with critical regulatory proteins, including E2F family transcription factors, c-Abl tyrosine kinase, and proteins containing a conserved LXCXE motif (1,2). In quiescent G_0 phase cells, active Rb proteins are hypophosphorylated and bind to E2F transcription factors to repress transcription and inhibit cell cycle progression (1,2). Upon growth factor induction of quiescent cells, Rb proteins become hyperphosphorylated and inactivated by G1-phase cyclinD-cdk4/6, G1/S-phase cyclin E-cdk2, and G1/S-phase cyclin A-cdk2 complexes (1,2). Hyperphosphorylation of Rb proteins results in a loss of E2F binding and allows for transcriptional activation and cell cycle progression (1,2). In addition to regulating the cell cycle, Rb proteins regulate chromosome stability, induction, and maintenance of senescence, apoptosis, cellular differentiation, and angiogenesis (3). Retinoblastoma-like protein 2 (RBL2, p130) is the most predominant and active Rb family member found in quiescent cells. In these cells, RBL2 interacts with E2F4 and E2F5 to recruit the DP, RB-like, E2F, and MuvB protein (DREAM) complex to E2F target genes to repress transcription of multiple genes required for progression into S phase and mitosis (4-6). Hypophosphorylation of RBL2 during cellular senescence is required for maintenance of senescent cells (7,8). **Background References** 1. Du, W. and Pogoriler, J. (2006) Oncogene 25, 5190-200. 2. Giacinti, C. and Giordano, A. (2006) Oncogene 25, 5220-7. 3. Indovina, P. et al. (2013) J Cell Physiol 228, 525-35. 4. Moberg, K. et al. (1996) Mol Cell Biol 16, 1436-49. 5. Takahashi, Y. et al. (2000) Genes Dev 14, 804-16. 6. Smith, E.J. et al. (1996) Mol Cell Biol 16, 6965-76. 7. Kapić, A. et al. (2006) Cell Death Differ 13, 324-34. 8. Helmbold, H. et al. (2009) Oncogene 28, 3456-67. Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot). Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween [®] 20 at 4°C with gentle shaking, overnight. Applications Key W: Western Blotting IP: Immunoprecipitation ChIP: Chromatin IP

Cross-Reactivity Key	H: Human M: Mouse R: Rat Mk: Monkey
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
	SignalSilence is a registered trademark of Cell Signaling Technology, Inc.
	SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.
	XP is a registered trademark of Cell Signaling Technology, Inc.
	All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.