FEZ1 Antibody	T C	Cell Signaling		
	Orders:	877-616-CELL (2355) orders@cellsignal.com		
2	Support:	877-678-TECH (8324)		
#14235	Web:	info@cellsignal.com cellsignal.com		
7	3 Trask Lane Danvers Mas	sachusetts 01923 USA		
For Research Use Only. Not for Use in Diagnostic Procedures.				

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 65	Source/Isotype: Rabbit	UniProt ID: #Q99689	Entrez-Gene Id: 9638
Product Usage Information	9	Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM soc 20°C. <i>Do not aliquot tl</i>		ö), 150 mM NaCl, 100 μg	/ml BSA and 50% gl	ycerol. Store at –
Specificity/Sen	sitivity	FEZ1 Antibody recogni	izes endogenous le	vels of total FEZ1 protei	n.	
Source / Purifi	cation		no terminus of hun	munizing animals with a nan FEZ1 protein. Antibo		
Background		The coiled-coil containing protein fasciculation and elongation protein zeta-1 (FEZ1) is expressed predominately in the brain and is the mammalian ortholog of the <i>C. elegans</i> protein UNC-76. It was identified independently in several interaction screens using distinct baits and was shown to play a role in neuronal differentiation and outgrowth, viral defense, centrosome organization, cytoskeletal signaling, and autophagy (reviewed in 1). It was originally identified as a binding partner and substrate for PKCζ and was found to induce the neuronal differentiation of PC-12 cells when co-expressed with active PKCζ (2). FEZ1 was also found to be an interacting partner with the schizophrenia-associated protein DISC1, which may suggest a role for FEZ1 in schizophrenia as well as other mental disorders (3,4). FEZ1 has also been shown to bind to several cytoskeletal proteins, including kinesins, tubulins, JIP1, NEK1, and CLASP2, which supports its role in neurite outgrowth, cargo transport along microtubules, and centrosomal organization (5-7). Additional research studies have shown that FEZ1 interacts with a viral agnoprotein and plays a role in viral defense, including during HIV-1 infection (8-10). Another screen identified FEZ1 as a binding partner for the ubiquitin ligase E4B and showed that FEZ1 can be regulated through polyubiquitination (11). Moreover, degradation of FEZ1 by the ubiquitination-proteasomal pathway through cdc20 provides a mechanism for FEZ1 in dendritic outgrowth (12). FEZ1 was also found to regulate autophagy through association with ULK1 and Beclin-1 complexes (13).				
Background R	eferences	2. Kuroda, S. et al. (199 3. Miyoshi, K. et al. (20 4. Kang, E. et al. (2011) 5. Fujita, T. et al. (2007 6. Blasius, T.L. et al. (20 7. Lanza, D.C. et al. (20 8. Suzuki, T. et al. (2009 9. Naghavi, M.H. et al.	99) J Cell Biol 144, 4 03) Mol Psychiatry) Neuron 72, 559-7) Biochem Biophys 007) J Cell Biol 176, 10) Mol Cell Bioche 5) J Biol Chem 280, (2005) Genes Dev 2009) Proc Natl Aca (2004) J Biol Chem 2 (2014) Cell Rep 7, 5	8, 685-94. I. <i>Res Commun</i> 361, 605- 11-7. em 338, 35-45. 24948-56. I9, 1105-15. I <i>d Sci U S A</i> 106, 14040-5 279, 53533-43. 552-64.	10.	
Species Reacti	vity	Species reactivity is de	termined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot E	Buffer	IMPORTANT: For west TBS, 0.1% Tween® 20		membrane with diluted shaking, overnight.	primary antibody i	n 5% w/v BSA, 1X
Applications K	ey	W: Western Blotting				
Cross-Reactivi	ty Key	H: Human M: Mouse F	R: Rat			

Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
	All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.