

Phospho-SQSTM1/p62 (Ser403) Antibody

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M	Endogenous	62	Rabbit	#Q13501	8878

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-SQSTM1/p62 (Ser403) Antibody recognizes endogenous levels of SQSTM1/p62 protein only when phosphorylated at Ser403.

Species predicted to react based on 100% sequence homology

Rat

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser403 of human SQSTM1/p62 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Sequestosome 1 (SQSTM1, p62) is a ubiquitin binding protein involved in cell signaling, oxidative stress, and autophagy (1-4). It was first identified as a protein that binds to the SH2 domain of p56Lck (5) and independently found to interact with PKCζ (6,7). SQSTM1 was subsequently found to interact with ubiquitin, providing a scaffold for several signaling proteins and triggering degradation of proteins through the proteasome or lysosome (8). Interaction between SQSTM1 and TRAF6 leads to the K63-linked polyubiquitination of TRAF6 and subsequent activation of the NF-κB pathway (9). Protein aggregates formed by SQSTM1 can be degraded by the autophagosome (4,10,11). SQSTM1 binds autophagosomal membrane protein LC3/Atg8, bringing SQSTM1-containing protein aggregates to the autophagosome (12). Lysosomal degradation of autophagosomes leads to a decrease in SQSTM1 levels during autophagy; conversely, autophagy inhibitors stabilize SQSTM1 levels. Studies have demonstrated a link between SQSTM1 and oxidative stress. SQSTM1 interacts with KEAP1, which is a cytoplasmic inhibitor of NRF2, a key transcription factor involved in cellular responses to oxidative stress (3). Thus, accumulation of SQSTM1 can lead to an increase in NRF2 activity. Phosphorylation of SQSTM1 at Ser403 increases its affinity for polyubiquitinated chains resulting in enhanced autophagic clearance (13,14). This site has been reported to be phosphorylated by casein kinase 2 (CK2) as well as by the innate immunity regulator TBK-1.

Background References

1. Kirkin, V. et al. (2009) *Mol Cell* 34, 259-69.
2. Seibenhener, M.L. et al. (2007) *FEBS Lett* 581, 175-9.
3. Komatsu, M. et al. (2010) *Nat Cell Biol* 12, 213-23.
4. Bjørkøy, G. et al. (2006) *Autophagy* 2, 138-9.
5. Joing, I. et al. (1996) *Proc Natl Acad Sci USA* 93, 5991-5.
6. Sanchez, P. et al. (1998) *Mol Cell Biol* 18, 3069-80.
7. Puls, A. et al. (1997) *Proc Natl Acad Sci USA* 94, 6191-6.
8. Vadlamudi, R.K. et al. (1996) *J Biol Chem* 271, 20235-7.
9. Wooten, M.W. et al. (2005) *J Biol Chem* 280, 35625-9.
10. Bjørkøy, G. et al. (2005) *J Cell Biol* 171, 603-14.
11. Komatsu, M. et al. (2007) *Cell* 131, 1149-63.
12. Pankiv, S. et al. (2007) *J Biol Chem* 282, 24131-45.
13. Matsumoto, G. et al. (2011) *Mol Cell* 44, 279-89.
14. Pilli, M. et al. (2012) *Immunity* 37, 223-34.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key**W:** Western Blotting**Cross-Reactivity Key****H:** Human **M:** Mouse**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.