

#14412 Store at -20C

Ras (G12V Mutant Specific) (D2H12) Rabbit mAb



Cell Signaling
TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
WB, W-S	H	Endogenous	21	Rabbit IgG	#P01116	3845

Product Usage Information	<p>Application</p> <p>Western Blotting</p> <p>Simple Western™</p>	<p>Dilution</p> <p>1:1000</p> <p>1:10 - 1:50</p>
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. <i>Do not aliquot the antibody.</i>	
Specificity / Sensitivity	Ras (G12V Mutant Specific) (D2H12) Rabbit mAb recognizes endogenous levels of Ras G12V mutant protein.	
Species predicted to react based on 100% sequence homology:	Mouse, Rat	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to G12V mutant sequence of human Ras protein.	
Background	<p>The 21 kDa guanine-nucleotide binding proteins (K-Ras, H-Ras, and N-Ras) cycle between active (GTP-bound) and inactive (GDP-bound) forms (1). Receptor tyrosine kinases and G protein-coupled receptors activate Ras, which then stimulates the Raf-MEK-MAPK pathway (2-4). GTPase-activating proteins (GAPs) normally facilitate the inactivation of Ras. However, research studies have shown that in 30% of human tumors, point mutations in Ras prevent the GAP-mediated inhibition of this pathway (5). The most common oncogenic Ras mutation found in tumors is Gly12 to Asp12 (G12D), which prevents Ras inactivation, possibly by increasing the overall rigidity of the protein (5,6).</p> <p>Additional oncogenic mutations of Ras have been observed at varying frequencies at codons 12, 13, and 61. The Gly12 to Val12 (G12V) mutation has been detected in a number of different cancers, including colorectal and thyroid cancer (7,8).</p>	
Background References	<ol style="list-style-type: none"> Boguski, M.S. and McCormick, F. (1993) <i>Nature</i> 366, 643-54. Avruch, J. et al. (1994) <i>Trends Biochem Sci</i> 19, 279-83. Buday, L. and Downward, J. (1993) <i>Cell</i> 73, 611-20. Huang, D.C. et al. (1993) <i>Mol Cell Biol</i> 13, 2420-31. Bos, J.L. (1989) <i>Cancer Res</i> 49, 4682-9. Ma, J. and Karplus, M. (1997) <i>J Mol Biol</i> 274, 114-31. Prior, I.A. et al. (2012) <i>Cancer Res</i> 72, 2457-67. Winder, T. et al. (2009) <i>Oncol Rep</i> 21, 1283-7. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
Applications Key	WB: Western Blotting W-S: Simple Western™
Cross-Reactivity Key	H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm: D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Sc: S. cerevisiae Ce: C. elegans Hr: horse GP: Guinea Pig Rab: rabbit All: all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.