**Revision** 1



## Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (197G2) Rabbit mAb (HRP Conjugate)



Orders: 877-616-CELL (2355) orders@cellsignal.com Support: 877-678-TECH (8324) Web: info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	<b>Reactivity:</b> H M R Mk Mi Dm Z Pg	Sensitivity: Endogenous	<b>MW (kDa):</b> 42, 44	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P27361, #P28482	<b>Entrez-Gene Id:</b> 5595, 5594
Product Usag Information	e	Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA, and 50% glycerol. Store at –20°C. Do not aliquot the antibodies.				
Specificity/Sensitivity		Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (197G2) Rabbit mAb (HRP Conjugate) detects endogenous levels of p44 and p42 MAP kinase (Erk1 and Erk2) when dually phosphorylated at Thr202 and Tyr204 of Erk1 (Thr185 and Tyr187 of Erk2), and singly phosphorylated at Tyr204. The antibody does not cross-react with the corresponding phosphorylated residues of either JNK/SAPK or p38 MAP kinase.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Thr202/Tyr204 of human p44 MAP kinase.				
Description		This Cell Signaling Technology antibody is conjugated to the carbohydrate groups of horseradish peroxidase (HRP) via its amine groups. The HRP conjugated antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (197G2) Rabbit mAb #4377.				
Background		Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs, such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli, including mitogens, growth factors, and cytokines (1-3), and research investigators consider it an important target in the diagnosis and treatment of cancer (4). Upon stimulation, a sequential three-part protein kinase cascade is initiated, consisting of a MAP kinase kinase kinase (MAPKK or MAP3K), a MAP kinase kinase (MAPKK or MAP3K), a MAP kinase (mAPKK or MAP2K), and a MAP kinase kinase mase (MAPKK or MAP3K), a MAP kinase (mAPKK or MAP2K), and a MAP kinase (MAPK). Multiple p44/42 MAP3Ks have been identified, including members of the Raf family, as well as Mos and Tpl2/COT. MEK1 and MEK2 are the primary MAPKKs in this pathway (5,6). MEK1 and MEK2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively. Several downstream targets of p44/42 have been identified, including p90RSK (7) and the transcription factor Elk-1 (8,9). p44/42 are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, known as DUSPs or MKPs (10), along with MEK inhibitors, such as U0126 and PD98059.				
Background References		<ol> <li>Roux, P.P. and Blenis, J. (2004) <i>Microbiol Mol Biol Rev</i> 68, 320-44.</li> <li>Baccarini, M. (2005) <i>FEBS Lett</i> 579, 3271-7.</li> <li>Meloche, S. and Pouysségur, J. (2007) <i>Oncogene</i> 26, 3227-39.</li> <li>Roberts, P.J. and Der, C.J. (2007) <i>Oncogene</i> 26, 3291-310.</li> <li>Rubinfeld, H. and Seger, R. (2005) <i>Mol Biotechnol</i> 31, 151-74.</li> <li>Murphy, L.O. and Blenis, J. (2006) <i>Trends Biochem Sci</i> 31, 268-75.</li> <li>Dalby, K.N. et al. (1998) <i>J Biol Chem</i> 273, 1496-505.</li> <li>Marais, R. et al. (1993) <i>Cell</i> 73, 381-93.</li> <li>Kortenjann, M. et al. (1994) <i>Mol Cell Biol</i> 14, 4815-24.</li> <li>Owens, D.M. and Keyse, S.M. (2007) <i>Oncogene</i> 26, 3203-13.</li> </ol>				
Species React	ivity	Species reactivity is de	etermined by testin	g in at least one appro	ved application (e.g., w	vestern blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat Mk: Monkey Mi: Mink Dm: D. melanogaster Z: Zebrafish Pg: Pig				

Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
	All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				
Limited Uses	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
	Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for reale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.				
Oudener 077 (46 CELL (2255)					

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com For Research Use Only. Not for Use in Diagnostic Procedures.