

Digoxigenin (D8Q9J) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: W	Reactivity: All	Sensitivity: Transfected Only	Source/Isotype: Rabbit IgG
Product Usage Information		Application Western Blotting	Dilution 1:20000
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.	
Specificity/Sensitivity		Digoxigenin (D8Q9J) Rabbit mAb recognizes digoxigenin conjugated molecules.	
Source / Purification		Monoclonal antibody is produced by immunizing animals with KLH-conjugated digoxigenin.	
Background		(1). This small hapten mole glycans. Antibodies target	only immunogenic steroid derived from "foxglove" plants of the genus <i>Digitalis</i> ecule can be covalently linked to proteins, nucleic acids, compounds, and sed against the DIG hapten display high binding affinity to DIG-conjugated d in molecular biology assays, including <i>in situ</i> hybridization, western blot, (2-4).
Background References		1. Cardwell, H.M. and Smith, S. (1953) <i>Experientia</i> 9, 367-8. 2. Kessler, C. (1991) <i>Mol Cell Probes</i> 5, 161-205. 3. Tian, F. et al. (2010) <i>Autophagy</i> 6, 1107-14. 4. Lin, L. et al. (2011) <i>J Virol Methods</i> 171, 91-7.	
Species Reactivit	y	Species reactivity is deterr	nined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key		W: Western Blotting	
Cross-Reactivity Key		All: All Species Expected	
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.	

more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c)

not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.