YAP (D8H1X) XP[®] Rabbit mAb (PE Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: FC-FP	Reactivity: H M R Hm Mk	Sensitivity: Endogenous	Source/Isotype: Rabbit IgG	UniProt ID: #P46937	Entrez-Gene Id: 10413
Product Usage Information		Application Flow Cytometry (Fixed/P	ermeabilized)		Dilution 1:50
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibodies. Protect from light. Do not freeze.			
Specificity/Sensitivity		YAP (D8H1X) XP [®] Rabbit mAb (PE Conjugate) recognizes endogenous levels of total YAP protein.			
Species predicted to react based on 100% sequence homology		Bovine, Horse, Guinea Pig			
Source / Purification		Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the carboxy terminus of human YAP protein. The epitope corresponds to a region surrounding Pro435 of human YAP isoform 1. This sequence region is 100% conserved among all known isoforms of human YAP protein.			
Description			alysis in human cells. The	antibody is expecte	n (PE) and tested in-house for d to exhibit the same species 1074.
Background		domain of Yes. It also bir In addition to the SH3 bir WW domains (2-4). While specific subcellular compactivator by virtue of its PEBP2 and other transcr widely recognized as a c conserved role in regula Ser109, Ser127) by LATS it is sequestered through	nds to other SH3 domain nding motif, YAP contain in initial studies of YAP all partments, subsequent sow domain interacting with a factors (5). In its call the factors of the Hi ting tissue growth and out kinases promotes YAP transsociation with 14-3-3 and for subsequent phosplice.	containing proteins a PDZ interaction mediated towards a rotudies showed that Youth the PY motif (PP) apacity as a transcrip ppo Pathway, which gan size (6-8). Phospanslocation from the proteins (7-9). These	oility to associate with the SH3 such as Nck, Crk, Src, and Abl (1). notif, a coiled-coil domain, and ole in anchoring and targeting to AP is a transcriptional co-cky) of the transcription factor tional co-activator, YAP is now plays a fundamental and widely shorylation at multiple sites (e.g., nucleus to the cytoplasm, where LATS-driven phosphorylation in an adjacent phosphodegron,
Background References		1. Sudol, M. (1994) <i>Oncogene</i> 9, 2145-52. 2. Mohler, P.J. et al. (1999) <i>J Cell Biol</i> 147, 879-90. 3. Espanel, X. and Sudol, M. (2001) <i>J Biol Chem</i> 276, 14514-23. 4. Sudol, M. et al. (1995) <i>FEBS Lett</i> 369, 67-71. 5. Yagi, R. et al. (1999) <i>EMBO J</i> 18, 2551-62. 6. Dong, J. et al. (2007) <i>Cell</i> 130, 1120-33. 7. Zhao, B. et al. (2010) <i>Genes Dev</i> 24, 862-74. 8. Zhao, B. et al. (2007) <i>Genes Dev</i> 21, 2747-61. 9. Yu, F.X. et al. (2012) <i>Cell</i> 150, 780-91. 10. Zhao, B. et al. (2010) <i>Genes Dev</i> 24, 72-85.			

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key H: Human M: Mouse R: Rat Hm: Hamster Mk: Monkey

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.