-20C

Store at

14752

#١





Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 95	Source/Isotype: Rabbit IgG	UniProt ID: #O43526	Entrez-Gene Id: 3785		
Product Usage Information		ApplicationDilutionWestern Blotting1:1000Immunoprecipitation1:50						
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				ol and less than		
Specificity/Sensitivity		Kv7.2 (D9L5S) Rabbit mAb recognizes endogenous levels of total Kv7.2 protein. This antibody does not cross-react with Kv7.3 protein.						
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human Kv7.2 protein.						
Background		The voltage gated potassium channel Kv7.2 (KCNQ2) associates with its family member Kv7.3 (KCNQ3) to form an M-channel that is involved in synaptic input response and sub-threshold excitability of neurons (1). This heteromeric channel generates the M-current, a slowly activating and deactivating potassium conductance that determines the neuronal excitability (2,3). Expression of these two M-channel proteins is mainly seen within the central nervous system, with both Kv7.2 and Kv7.3 expressed post-synaptically in the human cortex and hippocampus (4). The calcium-binding protein calmodulin binds two separate sites in Kv7.2 to influence exit of the channel protein from the endoplasmic reticulum and translocation to the plasma membrane (5). Mutations in the corresponding <i>KCNQ2</i> gene cause benign familial neonatal seizures-1 (BFNS1), an autosomal dominant form of epilepsy characterized by seizure clusters closely following birth (6,7).						
Background Ro	eferences	1. Stewart, A.P. et al. (2012) <i>J Biol Chem</i> 287, 11870-7. 2. Wang, H.S. et al. (1998) <i>Science</i> 282, 1890-3. 3. Smith, J.S. et al. (2001) <i>J Neurosci</i> 21, 1096-103. 4. Cooper, E.C. et al. (2000) <i>Proc Natl Acad Sci U S A</i> 97, 4914-9. 5. Alaimo, A. et al. (2013) <i>J Cell Sci</i> 126, 244-53. 6. Miraglia del Giudice, E. et al. (2000) <i>Eur J Hum Genet</i> 8, 994-7. 7. Dedek, K. et al. (2001) <i>Proc Natl Acad Sci U S A</i> 98, 12272-7.						
Species Reacti	vity	Species reactivity is det	termined by testin	g in at least one approve	ed application (e.g.,	western blot).		
Western Blot E	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				ו 5% w/v BSA, 1X		
Applications K	ey	W: Western Blotting IP: Immunoprecipitation						
Cross-Reactivit	ty Key	H: Human M: Mouse R: Rat						
Trademarks and Patents Cell Signaling Technology is a tradema				of Cell Signaling Technology, Inc.				
		XP is a registered trademark of Cell Signaling Technology, Inc.						
		All other trademarks an more information.	re the property of	their respective owners.	Visit cellsignal.com	/trademarks for		
Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized rep the following terms apply to Products provided by CST, its affiliates or its distributo terms and conditions that are in addition to, or different from, those contained her separately accepted in writing by a legally authorized representative of CST, are reju					s. Any Customer's in, unless			

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.