

Store at
-20C
#14783**VANGL1 (D1J7X) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Hm Mk	Endogenous	60	Rabbit IgG	#Q8TAA9	81839

Product Usage Information

Application
Western Blotting

Dilution
1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

VANGL1 (D1J7X) Rabbit mAb recognizes endogenous levels of total VANGL1 protein. This antibody does not cross-react with VANGL2 protein.

Species predicted to react based on 100% sequence homology

Chicken, Bovine, Pig, Horse

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu309 of human VANGL1 protein.

Background

Van Gogh-like proteins (VANGL1, VANGL2) are human orthologs of *Drosophila* Van Gogh (Vang/Stbm), a multi-pass transmembrane protein that is required to establish cell polarity in embryonic eyes, legs, and bristles (1,2). As in *Drosophila*, mammalian VANGL proteins are core components of the planar cell polarity (PCP) pathway that promotes asymmetric orientation of cells across a planar surface, and drives convergence-extension movements that are critical for tissue morphogenesis (3). Mutations in the human *VANGL1* gene have been identified in patients diagnosed with neural tube defects (e.g., spina bifida), providing evidence that VANGL1 plays a role in human embryonic morphogenesis (4,5). These findings are supported by genetic studies in mice, where mutations in both *Vangl1* and *Vangl2* result in neural tube defects (6,7). A possible role for VANGL in tumor progression is suggested by an increased expression of *VANGL1* mRNA in breast cancer patients with an elevated risk of relapse (8).

Background References

1. Wolff, T. and Rubin, G.M. (1998) *Development* 125, 1149-59.
2. Bastock, R. et al. (2003) *Development* 130, 3007-14.
3. Katoh, M. (2005) *Oncol Rep* 14, 1583-8.
4. Kibar, Z. et al. (2007) *N Engl J Med* 356, 1432-7.
5. Merello, E. et al. (2015) *Birth Defects Res A Clin Mol Teratol* 103, 51-61.
6. Iliescu, A. et al. (2011) *Biochemistry* 50, 795-804.
7. Chen, B. et al. (2013) *Genet Mol Res* 12, 3157-65.
8. Anastas, J.N. et al. (2012) *Oncogene* 31, 3696-708.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Hm:** Hamster **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.