

Store at
-20C
#14824**TRF2 (P449) Antibody**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H Mk	Endogenous	65	Rabbit	#Q15554	7014

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

TRF2 (P449) Antibody recognizes endogenous levels of total TRF2 protein. This antibody also cross-reacts with an unidentified protein of 135 kDa.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro449 of human TRF2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Telomeres, the linear ends of chromosomes, are organized into T-loops to prevent them from being recognized by the cell as DNA double stranded breaks (DSBs) (1). The telomeric repeat binding factor proteins TRF1 and TRF2 bind to double-stranded telomeres to allow formation of T-loops (2). A large number of proteins involved in the DNA damage response are found physically associated with TRF2 within telomeres (3). Interestingly, TRF2 can transiently localize to DNA damage-induced DSBs, but overexpression of TRF2 prevents ATM-dependent signaling (4). Phosphorylation of TRF2 at Ser323 has been reported *in vivo*, but no upstream kinase or role has been established for this phosphorylation site (5).

Background References

1. Karlseder, J. et al. (1999) *Science* 283, 1321-5.
2. Yoshimura, S.H. et al. (2004) *Genes Cells* 9, 205-18.
3. Colgin, L. and Reddel, R. (2004) *Curr Biol* 14, R901-2.
4. Bradshaw, P.S. et al. (2005) *Nat Genet* 37, 193-7.
5. Beausoleil, S.A. et al. (2004) *Proc Natl Acad Sci U S A* 101, 12130-5.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.