VIMP (D1D1M) Rabbit mAb



Orders:	877-616-CELL (2355) orders@cellsignal.com			
Support:	877-678-TECH (8324)			
Web:	info@cellsignal.com cellsignal.com			

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 21	Source/Isotype: Rabbit IgG	UniProt ID: #Q9BQE4	Entrez-Gene Id: 55829		
Product Usage Information		Application Western Blotting Immunoprecipitation			Dilution 1:1000 1:100			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Sen	Decificity/Sensitivity VIMP (D1D1M) Rabbit mAb recognizes endogenous levels of total VIMP protein.							
Source / Purific	cation	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys150 of human VIMP protein.						
Background		VCP-interacting membrane protein (VIMP, selenoprotein S) is a putative reductase and endoplasmic reticulum (ER)-resident protein involved in the ER-associated degradation (ERAD) pathway (1,2). Research studies indicate that VIMP may play a protective role against inflammation and reduce ER-stress (3). The VIMP protein is a single-pass, transmembrane protein that recruits the cytosolic p97/VCP AAA-ATPase and its cofactors, UFD1 and NPL4, to the ER membrane (4). An ER membrane complex containing Derlin-1 and VIMP forms a critical node in the ERAD machinery and links substrate recognition in the ER lumen with the retrotranslocation function of the p97/VCP AAA-ATPase in the cytosol (1,4). Polymorphisms in the corresponding <i>VIMP</i> gene are associated with spontaneous preterm births and cardiovascular disease risk (5,6) while other studies do not support a correspondence between <i>VIMP</i> polymorphisms and inflammatory disorders (7).						
Background Re	eferences	1. Lilley, B.N. and Ploegh, H.L. (2005) <i>Proc Natl Acad Sci U S A</i> 102, 14296-301. 2. Christensen, L.C. et al. (2012) <i>J Biol Chem</i> 287, 26388-99. 3. Fradejas, N. et al. (2011) <i>Glia</i> 59, 959-72. 4. Ye, Y. et al. (2004) <i>Nature</i> 429, 841-7. 5. Wang, Y. et al. (2013) <i>PLoS One</i> 8, e65657. 6. Cox, A.J. et al. (2013) <i>Acta Diabetol</i> 50, 391-9. 7. Martínez, A. et al. (2008) <i>BMC Genomics</i> 9, 329.						
Species Reactiv	vity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot B	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting IP: Immunoprecipitation						
Cross-Reactivit	су Кеу	H: Human						
Trademarks an	d Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		SignalSilence is a registered trademark of Cell Signaling Technology, Inc.						
		XP is a registered trademark of Cell Signaling Technology, Inc.						
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.						
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless						

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.