

TRPC6 (D3G1Q) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R	Endogenous	110	Rabbit IgG	#Q9Y210	7225

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

TRPC6 (D3G1Q) Rabbit mAb recognizes endogenous levels of total TRPC6 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human TRPC6 protein.

Background

Transient receptor potential (TRP) canonical 6 (TRPC6) belongs to the superfamily of TRP cation channels. The TRPC subfamily (TRPC1-7) is a group of calcium-permeable cation channels that mediates the increase in intracellular $[Ca^{2+}]$ following activation by G-protein-coupled receptors or receptor tyrosine kinases (1). TRPC6 is directly activated by diacylglycerol (DAG) (2). Various tissues express TRPC6, including brain, lung, kidney, ovary and small intestine. TRPC6 exerts a variety of biological functions in various tissues. In brain, TRPC6 plays important roles in synaptic plasticity, spatial cognition (3) and protects neurons from ischemic excitotoxicity (4). In kidney, TRPC6 is expressed in mesangial cells and podocytes of renal glomeruli (5) and regulates glomerular filtration (6). Mutations in TRPC6 gene cause autosomal dominant focal segmental glomerulosclerosis (7). In smooth muscle cells, TRPC6 mediates Na^+ influx followed by Ca^{2+} entry via Na^+/Ca^{2+} -exchanger (NCX) reversal which leads to contraction (8, 9).

Background References

1. Montell, C. (2005) *Sci STKE* 2005, re3.
2. Hofmann, T. et al. (1999) *Nature* 397, 259-63.
3. Liu, Y. et al. (2015) *Behav Brain Res* 294, 186-93.
4. Li, H. et al. (2012) *J Neurochem* 123, 1010-8.
5. Sours, S. et al. (2006) *Am J Physiol Renal Physiol* 290, F1507-15.
6. Dryer, S.E. and Reiser, J. (2010) *Am J Physiol Renal Physiol* 299, F689-701.
7. Reiser, J. et al. (2005) *Nat Genet* 37, 739-44.
8. Poburko, D. et al. (2007) *Circ Res* 101, 1030-8.
9. Dietrich, A. and Gudermann, T. (2014) *Handb Exp Pharmacol* 222, 157-88.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.