

BRF1/2 Antibody

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	40 to 50, 62	Rabbit	#P47974, #Q07352	678, 677

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

This antibody detects endogenous levels of total BRF1 and BRF2 proteins.

Species predicted to react based on 100% sequence homology

Chicken, Bovine

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the carboxy terminus of human BRF1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

ZFP36L1, also known as butyrate response factor-1 (BRF1), and ZFP36L2, also known as butyrate response factor-2 (BRF2), both belong to the TIS11 family of CCCH zinc-finger proteins (1). This family of proteins, which also includes tristetraprolin (TTP), bind to AU-rich elements (AREs) found in the 3'-untranslated regions of mRNAs and promote deadenylation and rapid degradation by the exosome (2,3). These proteins play a critical role in cell growth control by regulating the mRNA turnover of multiple cytokines, growth factors, and cell cycle regulators, including GM-CSF, TNFα, IL-2, IL-3, and IL-6 (4,5). Deregulated ARE-mRNA stability can contribute to both inflammation and oncogenic transformation (6-8). Insulin-induced stabilization of ARE-containing transcripts is mediated by Akt/PKB phosphorylation of ZFP36L1 at Ser92, which results in binding by 14-3-3 protein and inactivation of ZFP36L1 (9). ZFP36L1 and L2 have also been shown to promote cell quiescence in developing B lymphocytes, promoting VDJ recombination (10).

Background References

1. Varnum, B.C. et al. (1991) *Mol. Cell. Biol.* 11, 1754-1758.
2. Stoecklin, G. et al. (2002) *EMBO J.* 21, 4709-4718.
3. Lykke-Andersen, J. and Wagner, E. (2005) *Genes Dev.* 19, 351-361.
4. Stoecklin, G. et al. (2000) *Mol. Cell. Biol.* 20, 3753-3763.
5. Stoecklin, G. et al. (2001) *RNA* 7, 1578-1588.
6. Schuler, G.D. and Cole, M.D. (1988) *Cell* 55, 1115-1122.
7. Nair, A.P. et al. (1994) *Nature* 369, 239-242.
8. Carballo, E. et al. (1998) *Science* 281, 1001-1005.
9. Schmidlin, M. et al. (2004) *EMBO J.* 23, 4760-4769.
10. Galloway, A. et al. (2016) *Science* 352, 453-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.