

Store at -20C
#2152

WSTF Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 170, 185	Source/Isotype: Rabbit	UniProt ID: #Q9UIG0	Entrez-Gene Id: 9031
-------------------------------	--------------------------------	-----------------------------------	------------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

WSTF Antibody detects endogenous levels of total WSTF protein. The antibody does not cross-react with other related transcription factors.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the sequence of human WSTF (WBSCR9). Antibodies are purified by protein A and peptide affinity chromatography.

Background

The human WSTF gene is located within the common Williams Syndrome (WS) deletion area at chromosome 7q11.23. Several WSTF gene products have been detected with little difference in length of polypeptides (1-3). Functional motifs identified by sequence-homology searches include a PHD-type zinc finger motif followed by a bromodomain. Both motifs are found in many transcription factors, suggesting that WSTF may function as a transcription factor. A Drosophila gene (acf1) was cloned, which encodes two forms of Acf1 proteins with molecular weight 170 kDa and 185 kDa, respectively (4). It was demonstrated that Acf1 is structurally related to the human WSTF gene. Acf1 forms a complex with another protein, ISWI, and functions in the ATP-dependent catalysis of chromatin assembly (4).

Background References

1. Peoples, R.J. et al. (1998) *Cytogenet. Cell Genet.* 82, 238-246.
2. Lu, X. et al. (1998) *Genomics* 54, 241-249.
3. Jones, M.H. et al. (2000) *Genomics* 63, 40-45.
4. Ito, T. et al. (2000) *Genes Dev.* 13, 1529-1539.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.