e at -20C	Shh (C9C5) Rabbit mAb		Cell Signaling					
Stor		Orders:	877-616-CELL (2355) orders@cellsignal.com					
2		Support:	877-678-TECH (8324)					
207		Web:	info@cellsignal.com cellsignal.com					
#2		3 Trask Lane Danvers Mass	achusetts 01923 USA					

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H R	Sensitivity: Endogenous	MW (kDa): 19, (45 precursor)	Source/Isotype: Rabbit IgG	UniProt ID: #Q15465	Entrez-Gene Id: 6469			
Product Usage Information		Application Western Blotting			Dilution 1:1000				
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.							
Specificity/Sensitivity		Shh (C9C5) Rabbit mAb detects the endogenous levels of total Shh protein. This antibody does not cross-react with transfected IHH and DHH.							
Species predicted to react based on 100% sequence homology		Mouse, Chicken, Xenopus, Zebrafish							
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu53 of human Shh.							
Background		Hedgehog proteins (Hh) are secreted signaling proteins that play many roles during animal development. Aberrant Hh signaling activity can be associated with numerous birth defects and uncontrolled Hh pathway activation is linked to the development of several types of cancers (1-2). The three identified vertebrate Hh genes are Sonic (<i>Shh</i>), Indian (<i>Ihh</i>) and Desert (<i>Dhh</i>), all of which have distinct as well as overlapping roles (3-5). Hh proteins are synthesized as 45 kDa precursors that undergo auto-cleavage to generate a 19 kDa amino-terminal peptide (Hh-N) and a carboxy-terminal peptide (Hh-C). The Hh-N becomes covalently attached to a cholesterol molecule at its carboxy terminus and acetylated at its amino terminus. This doubly modified Hh-N peptide is released from cells and responsible for all known Hedgehog signaling activity (6).							
Background References		1. Ingham, P.W. and McMahon, A.P. (2001) <i>Genes Dev.</i> 15, 3059-3087. 2. McMahon, A.P. et al. (2003) <i>Curr. Top Dev. Biol.</i> 53, 1-114. 3. Zhang, X.M. et al. (2001) <i>Cell</i> 106, 781-792. 4. Adolphe, C. et al. (2004) <i>Development</i> 131, 5009-5019. 5. Pathi, S. et al. (2001) <i>Mech. Dev.</i> 106, 107-117. 6. Bijlsma, M.F. et al. (2004) <i>Bioessays</i> 26, 387-394.							
Species Reacti	vity	Species reactivity is	determined by testing	in at least one approve	ed application (e.g.,	western blot).			
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.							
Applications Key		W: Western Blotting							
Cross-Reactivity Key		H: Human R: Rat							
Trademarks a	Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
			U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.						
		All other trademarks more information.	s are the property of th	eir respective owners.	Visit cellsignal.com	/trademarks for			
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no							

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.