



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#2276

## Myc-Tag (9B11) Mouse mAb

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W, W-S, IP, IHC-P, IF-IC, FC-FP, ChIP	<b>Reactivity:</b> All	<b>Sensitivity:</b> Transfected Only	<b>Source/Isotype:</b> Mouse IgG2a kappa
<b>Product Usage Information</b>	For optimal ChIP results, use 5 µl of antibody and 10 µg of chromatin (approximately 4 x 10 <sup>6</sup> cells) per IP. This antibody has been validated using SimpleChIP <sup>®</sup> Enzymatic Chromatin IP Kits.		
	<b>Application</b>	<b>Dilution</b>	
	Western Blotting	1:1000	
	Simple Western™	1:10 - 1:50	
	Immunoprecipitation	1:250	
	Immunohistochemistry (Paraffin)	1:1500 - 1:6000	
	Immunofluorescence (Immunocytochemistry)	1:1000 - 1:2000	
	Flow Cytometry (Fixed/Permeabilized)	1:500 - 1:2000	
	Chromatin IP	1:50	
<b>Storage</b>	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.		
	For a carrier free (BSA and azide free) version of this product see product #58730.		
<b>Specificity/Sensitivity</b>	Myc-Tag (9B11) Mouse mAb detects exogenously expressed proteins containing the Myc epitope tag. This antibody recognizes the Myc tag fused to either the amino or carboxy terminus of targeted proteins in transfected cells. Myc-Tag (9B11) Mouse mAb detects exogenously expressed Myc-tagged proteins in cells expressed under a CMV promoter. Expression under other promoters has not been evaluated. The antibody may cross-react with c-myc protein. The antibody may weakly cross-react with a protein of unknown origin ~90kDa.		
<b>Source / Purification</b>	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues 410-419 of human c-Myc (EQKLISEEDL).		
<b>Background</b>	Epitope tags are useful for the labeling and detection of proteins using immunoblotting, immunoprecipitation, and immunostaining techniques. Because of their small size, they are unlikely to affect the tagged protein's biochemical properties.  The Myc epitope tag is widely used to detect expression of recombinant proteins in bacteria, yeast, insect and mammalian cell systems (1).		
<b>Background References</b>	1. Munro, S. and Pelham, H.R. (1984) <i>EMBO J.</i> 3, 3087-3093.		
<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).		
<b>Western Blot Buffer</b>	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.		
<b>Applications Key</b>	<b>W:</b> Western Blotting <b>W-S:</b> Simple Western™ <b>IP:</b> Immunoprecipitation <b>IHC-P:</b> Immunohistochemistry (Paraffin) <b>IF-IC:</b> Immunofluorescence (Immunocytochemistry) <b>FC-FP:</b> Flow Cytometry (Fixed/Permeabilized) <b>ChIP:</b> Chromatin IP		
<b>Cross-Reactivity Key</b>	<b>All:</b> All Species Expected		
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. SignalStain is a registered trademark of Cell Signaling Technology, Inc.		

SimpleChIP is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.