

p38γ MAPK Antibody



877-616-CELL (2355) Orders:

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 46	Source/Isotype: Rabbit	UniProt ID: #P53778	Entrez-Gene Id: 6300
Product Usage Information		Application Western Blotting Immunoprecipitation			Dilution 1:1000 1:100	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		p38γ MAPK Antibody detects endogenous levels of total p38γ MAPK protein. This antibody does not cross-react with the other p38 MAPK isoforms p38α, - β or - δ .				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the carboxy-terminal residues of human p38γ MAPK. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		p38 MAP kinase (MAPK), also called RK (1) or CSBP (2), is the mammalian orthologue of the yeast HOG kinase that participates in a signaling cascade controlling cellular responses to cytokines and stress (1-4). Four isoforms of p38 MAPK, p38α, β, γ (also known as Erk6 or SAPK3), and δ (also known as SAPK4) have been identified. Similar to the SAPK/JNK pathway, p38 MAPK is activated by a variety of cellular stresses, including osmotic shock, inflammatory cytokines, lipopolysaccharide (LPS), UV light, and growth factors (1-5). MKK3, MKK6, and SEK activate p38 MAPK by phosphorylation at Thr180 and Tyr182. Activated p38 MAPK has been shown to phosphorylate and activate MAPKAP kinase 2 (3) and to phosphorylate the transcription factors ATF-2 (5), Max (6), and MEF2 (5-8). SB203580 (4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)-imidazole) is a selective inhibitor of p38 MAPK. This compound inhibits the activation of MAPKAPK-2 by p38 MAPK and subsequent phosphorylation of HSP27 (9). SB203580 inhibits p38 MAPK catalytic activity by binding to the ATP-binding pocket, but does not inhibit phosphorylation of p38 MAPK by upstream kinases (10). Although there are many similarities between the four p38 isoforms, there are also some important differences that suggest that the various members may regulate specific functions, and the presence of multiple p38s may provide a mechanism for the generation of tissue-specific or stimulus-specific responses to the activation of the p38 signal transduction pathway (9,10).				
Background References		1. Rouse, J. et al. (1994) <i>Cell</i> 78, 1027-37. 2. Han, J. et al. (1994) <i>Science</i> 265, 808-11. 3. Lee, J.C. et al. (1994) <i>Nature</i> 372, 739-46. 4. Freshney, N.W. et al. (1994) <i>Cell</i> 78, 1039-49. 5. Raingeaud, J. et al. (1995) <i>J Biol Chem</i> 270, 7420-6. 6. Zervos, A.S. et al. (1995) <i>Proc Natl Acad Sci U S A</i> 92, 10531-4. 7. Zhao, M. et al. (1999) <i>Mol Cell Biol</i> 19, 21-30. 8. Yang, S.H. et al. (1999) <i>Mol Cell Biol</i> 19, 4028-38. 9. Cuenda, A. et al. (1995) <i>FEBS Lett</i> 364, 229-33. 10. Kumar, S. et al. (1999) <i>Biochem Biophys Res Commun</i> 263, 825-31. 11. Fearns, C. et al. (2000) <i>J. Leukoc. Biol.</i> 67, 705-711. 12. Hale, K.K. et al. (1999) <i>J. Immunol.</i> 162, 4246-4252.				

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key H: Human M: Mouse R: Rat Mk: Monkey

Trademarks and PatentsCell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.