



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

PAR-4 Antibody

Store at -20C
#2328

For Research Use Only. Not for Use in Diagnostic Procedures.

| | | | | | | |
|---|--------------------------------|-----------------------------------|------------------------|----------------------------------|-------------------------------|--------------------------------|
| Applications: W, IP, IF-IC, FC-FP | Reactivity: H M R Mk | Sensitivity: Endogenous | MW (kDa): 41 | Source/Isotype: Rabbit | UniProt ID: #Q961Z0 | Entrez-Gene Id: 5074 |
|---|--------------------------------|-----------------------------------|------------------------|----------------------------------|-------------------------------|--------------------------------|

Product Usage Information

Application

Western Blotting
Immunoprecipitation
Immunofluorescence (Immunocytochemistry)
Flow Cytometry (Fixed/Permeabilized)

Dilution

1:1000
1:100
1:100
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

PAR-4 Antibody detects endogenous levels of PAR-4 protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Arg220 of human PAR-4. Antibodies were purified by protein A and peptide affinity chromatography.

Background

PAR-4 (prostate apoptosis response-4) was identified as a protein that is upregulated in prostate tumor cells undergoing apoptosis (1). Additionally, in parallel studies PAR-4 was found in the yeast two-hybrid system to bind to the Wilms' tumor suppressor protein WT1 and may modulate WT1-mediated transcriptional activation (2). PAR-4 contains a leucine zipper domain and a death domain and has been implicated as an effector of apoptosis during tumorigenesis as well as in neurodegenerative disorders (3,4). PAR-4 is widely expressed in normal tissues but can be downregulated in some tumor types. The mechanism of PAR-4 mediated apoptosis regulation appears to be complex and dependent on the cellular context. Studies have indicated roles for PAR-4 in activation of the Fas-FADD-caspase-8 pathway as well as inhibition of the NF-κB pro-survival pathway (5-7). Its activity is likely to depend on the cellular context and post-translational modifications. For instance, phosphorylation of PAR-4 by Akt prevents its nuclear translocation thereby promoting cell survival (8). In contrast, phosphorylation of rat PAR-4 at T155 by PKA appears to positively regulate its apoptotic activity (9).

Background References

- Sells, S.F. et al. (1997) *Mol. Cell Biol.* 17, 3823-3832.
- Johnstone, R.W. et al. (1996) *Mol. Cell Biol.* 16, 6945-6956.
- Guo, Q. et al. (1998) *Nat. Med.* 4, 957-962.
- El-Guendy, N. and Rangnekar, V.M. (2003) *Exp. Cell Res.* 283, 51-66.
- Chakraborty, M. et al. (2001) *Cancer Res.* 61, 7255-7263.
- Díaz-Meco, M.T. et al. (1996) *Cell* 86, 777-786.
- Díaz-Meco, M.T. et al. (1999) *J. Biol. Chem.* 274, 19606-79612.
- Goswami, A. et al. (2005) *Mol. Cell* 20, 33-44.
- Gurumurthy, S. et al. (2005) *Mol. Cell Biol.* 25, 1146-1161.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.