

Revision 1

Store at  
-20C  
#24032**BCAR3 Antibody**
**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H	Endogenous	95	Rabbit	#O75815	8412

**Product Usage Information****Application**Western Blotting  
Immunoprecipitation**Dilution**1:1000  
1:100**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

BCAR3 antibody recognizes endogenous levels of total BCAR3 protein.

**Source / Purification**

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro526 of human BCAR3 protein. Antibodies are purified by protein A and peptide affinity chromatography.

**Background**

BCAR3 is a member of the novel SH2-containing protein (NSP) family (1). It was identified as a gene product involved in anti-estrogen resistance in the context of breast cancer (2). Like other members of this family, BCAR3 has been shown to interact with the family member, CAS. The C terminal Cdc25 homology domain of BCAR3 interacts tightly with the FAT domain of p130Cas (3) and promotes the association of p130cas with Src kinase (4) to activate related signaling pathways. Overexpression of BCAR3 leads to the activation of a wide range of downstream signaling proteins including PI3K, rac, PAK1, and cyclin D1 (5-7). The main role of BCAR3 is to promote cell motility and regulate cytoskeletal remodeling and adhesion through its effect on p130cas and Src kinase (8-10). BCAR3 also has been implicated in playing an inhibitory role on TGF-β/SMAD signaling, which is associated with favorable disease outcomes (11).

**Background References**

1. Near, R.I. et al. (2007) *J Cell Physiol* 212, 655-65.
2. van Agthoven, T. et al. (1998) *EMBO J* 17, 2799-808.
3. Mace, P.D. et al. (2011) *Nat Struct Mol Biol* 18, 1381-7.
4. Makkinje, A. et al. (2012) *J Biol Chem* 287, 27703-14.
5. Cai, D. et al. (2003) *Cancer Res* 63, 6802-8.
6. Felekis, K.N. et al. (2005) *Mol Cancer Res* 3, 32-41.
7. Cai, D. et al. (2003) *J Immunol* 170, 969-78.
8. Schuh, N.R. et al. (2010) *J Biol Chem* 285, 2309-17.
9. Wilson, A.L. et al. (2013) *PLoS One* 8, e65678.
10. Makkinje, A. et al. (2009) *Cell Signal* 21, 1423-35.
11. Guo, J. et al. (2014) *Breast Cancer Res* 16, 476.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

**Applications Key****W:** Western Blotting **IP:** Immunoprecipitation**Cross-Reactivity Key****H:** Human**Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.**Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

**Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com**  
For Research Use Only. Not for Use in Diagnostic Procedures.