

Store at
-20C
#24083**OGT (D1D8Q) Rabbit mAb**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 110	Source/Isotype: Rabbit IgG	UniProt ID: #O15294	Entrez-Gene Id: 8473
-------------------------------	--------------------------------	-----------------------------------	-------------------------	--------------------------------------	-------------------------------	--------------------------------

Product Usage Information	Application Western Blotting Immunoprecipitation	Dilution 1:1000 1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity/Sensitivity	OGT (D1D8Q) Rabbit mAb recognizes endogenous levels of total OGT protein.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro546 of human OGT protein.	
Background	O-GlcNAcylation is a post-translational modification where β-D-N-acetylglucosamine (GlcNAc) is covalently linked to cytoplasmic and nuclear proteins at serine or threonine residues (1,2). This modification is important in many cellular processes including metabolism, cell growth and morphogenesis, apoptosis, and transcription (2,3), and research studies have implicated this modification in cancer (1). The reversible protein modification by O-GlcNAc, which has been suggested to be a nutrient and stress sensor, is catalyzed by two highly conserved enzymes, O-GlcNAc transferase (OGT) and O-GlcNAcase (OGA) (4).	
Background References	<ol style="list-style-type: none"> 1. Comer, F.I. et al. (2001) <i>Anal Biochem</i> 293, 169-77. 2. Slawson, C. and Hart, G.W. (2011) <i>Nat Rev Cancer</i> 11, 678-84. 3. Capotosti, F. et al. (2011) <i>Cell</i> 144, 376-88. 4. Hart, G.W. et al. (2007) <i>Nature</i> 446, 1017-22. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key	W: Western Blotting IP: Immunoprecipitation	
Cross-Reactivity Key	H: Human M: Mouse R: Rat Mk: Monkey	
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.	
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party,</p>	

whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.