



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C  
#2410

## Clathrin Heavy Chain (P1663) Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IF-IC	<b>Reactivity:</b> H M R Mk B	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 190	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #Q00610	<b>Entrez-Gene Id:</b> 1213
----------------------------------	----------------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	--------------------------------

### Product Usage Information

#### Application

Western Blotting  
Immunofluorescence (Immunocytochemistry)

#### Dilution

1:1000  
1:100

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

Clathrin Heavy Chain (P1663) Antibody detects endogenous levels of total Clathrin protein.

### Species predicted to react based on 100% sequence homology

Pig

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro1663 of human Clathrin. Antibodies are purified by protein A and peptide affinity chromatography

### Background

Clathrin-coated vesicles provide for the intracellular transport of cargo proteins following endocytosis and during multiple vesicle trafficking pathways. Vesicles form at specialized areas of the cell membrane where clathrin and associated proteins form clathrin-coated pits. Invagination of these cell membrane-associated pits internalizes proteins and forms an intracellular clathrin-coated vesicle (1,2). Clathrin is the most abundant protein in these vesicles and is present as a basic assembly unit called a triskelion. Each clathrin triskelion is composed of three clathrin heavy chains and three clathrin light chains. Clathrin heavy chain proteins are composed of several functional domains, including a carboxy-terminal region that permits interaction with other heavy chain proteins within a triskelion, and a globular amino-terminal region that associates with other vesicle proteins (2). Adaptor proteins, such as AP2, epsin and EPS15, are responsible for the recruitment of vesicle proteins to sites of pit formation and the assembly of the clathrin-coated vesicle. Following vesicle invagination, the GTPase dynamin constricts the neck of the nascent vesicle to complete formation of the free, cytosolic vesicle (3,4).

### Background References

- Rodriguez-Boulan, E. et al. (2005) *Nat. Rev. Mol. Cell Biol.* 6, 233-247.
- Mousavi, S.A. et al. (2004) *Biochem. J.* 377, 1-16.
- Rappoport, J.Z. et al. (2004) *Traffic* 5, 327-237.
- Brett, T.J. and Traub, L.M. (2006) *Curr. Opin. Cell Biol.* 18, 395-406.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **IF-IC:** Immunofluorescence (Immunocytochemistry)

### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey **B:** Bovine

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.