



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#2443

PPAR γ (81B8) Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

| | | | | | | |
|---|---------------------------|-----------------------------------|----------------------------|----------------------------------|-------------------------------|--------------------------------|
| Applications: W, IP, IF-IC, ChIP, ChIP-seq | Reactivity: H M | Sensitivity: Endogenous | MW (kDa): 53, 57 | Source/Isotype: Rabbit | UniProt ID: #P37231 | Entrez-Gene Id: 5468 |
|---|---------------------------|-----------------------------------|----------------------------|----------------------------------|-------------------------------|--------------------------------|

Product Usage Information

For optimal ChIP and ChIP-seq results, use 5 μ l of antibody and 10 μ g of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP[®] Enzymatic Chromatin IP Kits.

| Application | Dilution |
|--|---------------|
| Western Blotting | 1:1000 |
| Immunoprecipitation | 1:50 |
| Immunofluorescence (Immunocytochemistry) | 1:100 - 1:200 |
| Chromatin IP | 1:100 |
| Chromatin IP-seq | 1:100 |

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

PPAR γ (81B8) Rabbit mAb detects endogenous levels of total PPAR γ protein.

Species predicted to react based on 100% sequence homology

Rat

Source / Purification

PPAR γ (81B8) Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding His494 of human PPAR γ .

Background

Peroxisome proliferator-activated receptor γ (PPAR γ) is a member of the ligand-activated nuclear receptor superfamily and functions as a transcriptional activator (1). PPAR γ is preferentially expressed in adipocytes as well as in vascular smooth muscle cells and macrophage (2). Besides its role in mediating adipogenesis and lipid metabolism (2), PPAR γ also modulates insulin sensitivity, cell proliferation and inflammation (3). PPAR γ transcriptional activity is inhibited by MAP kinase phosphorylation of PPAR γ at Ser84 (4,5).

Background References

1. Tontonoz, P. et al. (1995) *Curr. Opin. Genet. Dev.* 5, 571-576.
2. Rosen, E.D. et al. (1999) *Mol. Cell* 4, 611-617.
3. Murphy, G.J. and Holder, J.C. (2000) *Trends Pharmacol. Sci.* 21, 469-474.
4. Camp, H.S. and Tafuri, S.R. (1997) *J. Biol. Chem.* 272, 10811-10816.
5. Adams, M. et al. (1997) *J. Biol. Chem.* 272, 5128-5132.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry)
ChIP: Chromatin IP **ChIP-seq:** Chromatin IP-seq

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.