Store at -20C

## eIF5 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	58	Rabbit	#P55010	1983
Product Usage Information		<b>Application</b> Western Blotting			<b>Dilution</b> 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		eIF5 Antibody detects endogenous levels of total eIF5 protein.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to a sequence surrounding Val417 of human eIF5. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		Eukaryotic translation initiation factor 5 (eIF5) is crucial for the assembly of translation initiation complex and plays an important role in protein synthesis (1). eIF5 interacts with the 43S initiation complex to stimulate hydrolysis of GTP bound to eIF2 (1-3). Studies suggest that eIF5 functions as the GTPase-activating protein (GAP) in the hydrolysis of GTP-bound eIF2 (4,5). This hydrolysis leads to the release of initiation factors from the 40S ribosomal subunit, which is a necessary step in the formation of the 80S initiation complex (1).				
Background References		1. Das, S. and Maitra, U. (2001) <i>Prog Nucleic Acid Res Mol Biol</i> 70, 207-31. 2. Chakrabarti, A. and Maitra, U. (1991) <i>J Biol Chem</i> 266, 14039-45. 3. Asano, K. et al. (2000) <i>Genes Dev</i> 14, 2534-46. 4. Das, S. et al. (2001) <i>J Biol Chem</i> 276, 6720-6. 5. Paulin, F.E. et al. (2001) <i>Curr Biol</i> 11, 55-9.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat Mk: Monkey				
<b>Trademarks and Patents</b>		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks a more information.	re the property of	their respective owners.	Visit cellsignal.com	ı/trademarks for
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
		approved, cleared, or purpose. Customer sh any manner that confl	licensed by the FDA all not use any Pro icts with its labelin	se Only or a similar labe to or other regulatory for duct for any diagnostic of g statement. Products greearch and development	eign or domestic er or therapeutic purp old or licensed by C	ntity, for any lose, or otherwise in IST are provided for

Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party,

whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.