

*475*59

EGR3 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 50	Source/Isotype: Rabbit	UniProt ID: #Q06889	Entrez-Gene Id: 1960	
Product Usage Information	•	Application Western Blotting			Dilution 1:1000		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity/Sensitivity		EGR3 Antibody recognizes endogenous levels of total EGR3 protein.					
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Phe160 of human EGR3 protein. Antibodies are purified by protein A and peptide affinity chromatography.					
Background		Early growth response 3 (EGR3) is a zinc finger transcription factor and one of four members of the early growth response family (1). Members of this family are basally expressed in the brain (2) and are immediate early response genes that are important for the induction of cellular programs of differentiation, proliferation, and cell death in response to environmental stimuli (1). EGR3 plays important roles in cellular growth and neuronal development (2) and an essential role in learning and memory processing of both short- and long-term hippocampus-dependent memory; it also mediates adaptation to stress and novelty (3). Research studies show that a lack of EGR3 results in abnormal neuronal and T cell development (4).					
Background References		1. Kumbrink, J. et al. (2010) <i>J Cell Biochem</i> 111, 207-17. 2. Kim, S.H. et al. (2010) <i>Am J Med Genet B Neuropsychiatr Genet</i> 153B, 1355-60. 3. Zhang, R. et al. (2012) <i>PLoS One</i> 7, e30237. 4. Suehiro, J. et al. (2010) <i>Blood</i> 115, 2520-32.					
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Key		W: Western Blotting					
Cross-Reactivity Key		H: Human M: Mouse R: Rat					
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.					
		All other trademarks a more information.	are the property of t	heir respective owners.	Visit cellsignal.com	/trademarks for	
Limited Uses		the following terms ar terms and conditions	Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.				
		Products are labeled with For Research Use Only or a similar labeling statement and have not been					

approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.