

## **PP1α Antibody**



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	<b>Reactivity:</b> H M R Mk Dm	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 38	<b>Source/Isotype:</b> Rabbit	UniProt ID: #P62136	Entrez-Gene Id: 5499
Product Usage Information		<b>Application</b> Western Blotting			<b>Dilution</b> 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		PP1 $\alpha$ Antibody detects endogenous levels of total PP1 $\alpha$ protein. This antibody detects recombinant levels of PP1 $\gamma$ and does not cross-react with PP1 $\delta$ .				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the amino-terminal sequence of human PP1alpha. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		Type 1 protein phosphatase (PP1), a serine/threonine phosphatase, is highly conserved in eukaryotic cells. Four isoforms of PP1 have been characterized: PP1 $\alpha$ , PP1 $\delta$ , PP1 $\gamma$ 1 and PP1 $\gamma$ 2 (1). Involvement in cell cycle regulation is one of the biological functions of PP1 (1). It has been illustrated that PP1 dephosphorylates Rb and cdc25 during mitosis (2,3). A cell cycle-dependent phosphorylation at Thr320 of PP1 $\alpha$ by cdc2 kinase inhibits PP1 $\alpha$ activity (4).				
Background References		1. Rubin, E. et al. (1998) <i>Front Biosci</i> 3, D1209-19. 2. Durfee, T. et al. (1993) <i>Genes Dev</i> 7, 555-69. 3. Izumi, T. et al. (1992) <i>Mol Biol Cell</i> 3, 927-39. 4. Kwon, Y.G. et al. (1997) <i>Proc Natl Acad Sci U S A</i> 94, 2168-73.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat Mk: Monkey Dm: D. melanogaster				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless				

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any

commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.