



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#2701

Phospho-Zap-70 (Tyr319)/Syk (Tyr352) Antibody

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, W-S, IF-IC	H	Endogenous	70 Zap-70, 72 Syk	Rabbit	#P43403, #P43405	7535, 6850

Product Usage Information

Application

Western Blotting
Simple Western™
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:10 - 1:50
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Phospho-Zap-70 (Tyr319)/Syk (Tyr352) Antibody detects endogenous levels of Zap-70 only when phosphorylated at Tyr319. It cross-reacts with endogenous levels of Syk when phosphorylated at Tyr352.

Species predicted to react based on 100% sequence homology

Mouse, Rat, Hamster, Monkey, Chicken, Bovine, Dog, Pig, Horse

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr319 of human Zap-70. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The Syk family protein tyrosine kinase Zap-70 is expressed in T and NK cells and plays a critical role in mediating T cell activation in response to T cell receptor (TCR) engagement (1). Following TCR engagement, Zap-70 is rapidly phosphorylated on several tyrosine residues through autophosphorylation and transphosphorylation by the Src family tyrosine kinase Lck (2-6). Tyrosine phosphorylation correlates with increased Zap-70 kinase activity and downstream signaling events. Expression of Zap-70 is correlated with disease progression and survival in patients with chronic lymphocytic leukemia (7,8). Phosphorylation of Tyr319 is required for the assembly of a Zap-70-containing signaling complex that leads to the activation of the PLC-gamma1-dependent and Ras-dependent signaling cascades in antigen-stimulated T cells (5,6). The orthologous Tyr352 residue in Syk is also involved in the association with PLC-gamma1 (9).

Background References

1. Chu, D.H. et al. (1998) *Immunol Rev* 165, 167-80.
2. Iwashima, M. et al. (1994) *Science* 263, 1136-9.
3. Neumeister, E.N. et al. (1995) *Mol Cell Biol* 15, 3171-8.
4. Chan, A.C. et al. (1995) *EMBO J* 14, 2499-508.
5. Williams, B.L. et al. (1999) *EMBO J* 18, 1832-44.
6. Di Bartolo, V. et al. (1999) *J Biol Chem* 274, 6285-94.
7. Wiestner, A. et al. (2003) *Blood* 101, 4944-51.
8. Crespo, M. et al. (2003) *N Engl J Med* 348, 1764-75.
9. Law, C.L. et al. (1996) *Mol Cell Biol* 16, 1305-15.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **W-S:** Simple Western™ **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.