NEDD8 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

877-678-TECH (8324) Support:

info@cellsignal.com cellsignal.com Web:

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IHC-P	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 9	Source/Isotype: Rabbit	UniProt ID: #Q15843	Entrez-Gene Id: 4738
Product Usage Information		Application Western Blotting Immunoprecipitation Immunohistochemistry (Paraffin)			Dilution 1:1000 1:25 1:200	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		This antibody detects endogenous levels of both free and conjugated NEDD8 protein. The antibody does not cross-react with other ubiquitin family members, including ubiquitin, SUMO1, SUMO2, SUMO3 and ISG15.				
Species predicted to react based on 100% sequence homology		Xenopus, Zebrafish, Bo	ovine			
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to amino acids at the amino-terminus of human NEDD8 protein. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		Neural precursor cell-expressed developmentally downregulated protein 8 (NEDD8), also known as Rub1 (related to ubiquitin 1) in plants and yeast, is a member of the ubiquitin-like protein family (1,2). The covalent attachment of NEDD8 to target proteins, termed neddylation, is a reversible, multi-step process analogous to ubiquitination. NEDD8 is first synthesized in a precursor form with a carboxy-terminal extension peptide that is removed by either the UCH-L3 or NEDP1/DEN1 hydrolase protein to yield a mature NEDD8 protein (3,4). Mature NEDD8 is then covalently linked to target proteins via the carboxy-terminal glycine residue in a reaction catalyzed by the APP-BP1/Uba3 heterodimer complex and Ubc12 as the E1- and E2-like enzymes, respectively (5). An E3 ligase protein, Roc1/Rbx1, is also required for neddylation of the cullin proteins (6). Protein de-neddylation is catalyzed by a number of enzymes in the cell, including a "ubiquitin-specific" protease USP21, the NEDP1/DEN1 hydrolase and the COP9/signalosome (CSN) (7,8,9). In contrast to the ubiquitin pathway, the NEDD8 modification system acts on only a few substrates and does not appear to target proteins for degradation. Neddylation of cullin proteins activates the SCF (Skp1-Cullin-F-box) E3 ubiquitin ligase complex by promoting complex formation and enhancing the recruitment of the E2-ubiquitin intermediate (10). While NEDD8 modification of VHL is not required for ubiquitination of HIF1-α, it is required for fibronectin matrix assembly (11). Mdm2-dependent neddylation of p53 inhibits its transcriptional activity (12).				
Background References		1. Chiba, T. and Tanaka, K. (2004) <i>Curr. Protein Pept. Sci.</i> 5, 177-184. 2. Schwartz, D.C. and Hochstrasser, M. (2003) <i>Trends Biochem. Sci.</i> 28, 321-328. 3. Wada, H. et al. (1998) <i>Biochem. Biophys. Res. Commun.</i> 251, 688-692. 4. Hemelaar, J. et al. (2004) <i>Mol. Cell Biol.</i> 24, 84-95. 5. Osaka, F. et al. (1998) <i>Genes Dev.</i> 12, 2263-2268. 6. Kamura, T. et al. (1999) <i>Genes Dev.</i> 13, 2928-2933. 7. Gong, L. et al. (2000) <i>J. Biol. Chem.</i> 275, 14212-14216. 8. Mendoza, H.M. et al. (2003) <i>J. Biol. Chem.</i> 278, 25637-25643. 9. Lyapina, S. et al. (2001) <i>Science</i> 292, 1382-1385. 10. Kawakami, T. et al. (2001) <i>EMBO J.</i> 20, 4003-4012. 11. Stickle, N.H. et al. (2004) <i>Mol. Cell Biol.</i> 24, 3251-3261.				

12. Xirodimas, D.P. et al. (2004) Cell 118, 83-97.

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting IP: Immunoprecipitation IHC-P: Immunohistochemistry (Paraffin)

Cross-Reactivity Key H: Human M: Mouse R: Rat Mk: Monkey

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited UsesExcept as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless

separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.