:2839

SHIP2 (C76A7) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-IC, FC-FP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 160	Source/Isotype: Rabbit IgG	UniProt ID: #O15357	Entrez-Gene Id: 3636
Product Usage Information		Application Western Blotting Immunoprecipitation				Dilution 1:1000 1:50
		Immunofluorescence (Immunocytochemistry) Flow Cytometry (Fixed/Permeabilized)				1:25 1:50
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
		For a carrier free (BSA and azide free) version of this product see product #72173.				
Specificity/Sensitivity		SHIP2 (C76A7) Rabbit mAb detects endogenous levels of total SHIP2 protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala1083 of human SHIP2.				
Background		SH2-containing inositol phosphatase 1 (SHIP1) is a hematopoietic phosphatase that hydrolyzes phosphatidylinositol-3,4,5-triphosphate to phosphatidylinositol-3,4-bisphosphate (1). SHIP1 is a cytosolic phosphatase with an SH2 domain in its amino terminus and two NPXY Shc binding motifs in its carboxy terminus (1,2). Upon receptor cross-linking, SHIP is first recruited to the membrane junction through binding of its SH2 domain to the phospho-tyrosine in the ITIM motif (2), followed by tyrosine phosphorylation on the NPXY motif (2). The membrane relocalization and phosphorylation on the NPXY motif is essential for the regulatory function of SHIP1 (3-5). Its effect on calcium flux, cell survival, growth, cell cycle arrest, and apoptosis is mediated through the PI3K and Akt pathways (3-5). Tyr1022 is located in one of the NPXY motifs in SHIP1, and its phosphorylation is important for SHIP1 function (6).				
		SHIP2, a homolog of SHIP1, is highly expressed in heart, skeletal muscle and placenta (7). SHIP2 negatively regulates insulin signaling (8) and polymorphisms in SHIP2 have been linked to hyperglycemia (9). Recent studies also suggest SHIP2 as a therapeutic target for the treatment of both obesity and type 2 diabetes (10,11).				
Background Re	ferences	 Tridandapani, S. et al. (1997) Mol Cell Biol 17, 4305-11. Liu, L. et al. (1997) J Biol Chem 272, 8983-8. Malbec, O. et al. (2001) J Biol Chem 276, 30381-91. Carver, D.J. et al. (2000) Blood 96, 1449-56. Scharenberg, A.M. et al. (1998) EMBO J 17, 1961-72. Sattler, M. et al. (2001) J Biol Chem 276, 2451-8. Pesesse, X. et al. (1997) Biochem Biophys Res Commun 239, 697-700. Wada, T. et al. (2001) Mol Cell Biol 21, 1633-46. Ishida, S. et al. (2006) Pancreas 33, 63-7. Dyson, J.M. et al. (2005) Int J Biochem Cell Biol 37, 2260-5. Sasaoka, T. et al. (2006) Pharmacol Ther 112, 799-809. 				

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permashilized)

FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 7,429,487, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com

For Research Use Only. Not for Use in Diagnostic Procedures.